



Nicotinic Acetylcholine Receptors in HIV: Possible Roles During HAND and Inflammation

Coral M. Capó-Vélez^{1,2} · Manuel Delgado-Vélez^{1,2} · Carlos A. Báez-Pagán^{1,3} · José A. Lasalde-Dominicci^{1,2}

Received: 31 January 2018 / Accepted: 9 July 2018 / Published online: 14 July 2018
© The Author(s) 2018

Abstract

Infection with the human immunodeficiency virus (HIV) remains a threat to global health. Since its discovery, many efforts have been directed at understanding the mechanisms and consequences of infection. Although there have been substantial advances since the advent of antiretroviral therapy, there are still complications that significantly compromise the health of infected patients, particularly, chronic inflammation and HIV-associated neurocognitive disorders (HAND). In this review, a new perspective is addressed in the field of HIV, where the $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ -nAChR) is the protagonist. We comprehensively discuss the available evidence implicating $\alpha 7$ -nAChRs in the context of HIV and provide possible explanations about its role in HAND and inflammation in both the central nervous system and the periphery.

Keywords Nicotinic acetylcholine receptor · HIV · Gp120 · Inflammation · HAND

Introduction

Despite the success of the combined antiretroviral therapy (cART), infection with the human immunodeficiency virus type 1 (HIV-1) remains a global health threat. cART has helped prolong the lives of patients, improve their life quality, lessen the occurrence of the acquired immunodeficiency syndrome (AIDS), and decrease AIDS and non-AIDS-related complications; however, it is neither a cure (Spudich 2013) nor is it accessible to every person living with HIV (out of the approximately 36.7 million people infected with the virus, only 18.2 million have access to cART) (Joint United Nations Programme on HIV/AIDS (UNAIDS 2013)).

Besides AIDS, HIV infection results in a series of systematic physiological complications such as chronic inflammation and HIV-associated neurocognitive disorders (HAND), regardless of cART treatment (Appay and Sauce 2008; Heaton et al. 2011; Harezlak et al. 2011; Funderburg et al. 2014). These complications do not receive sufficient attention and continue to be a serious threat to the quality of life of patients living with the virus (Saylor et al. 2016; Althoff et al. 2016). Sadly, the mechanisms leading to either of these complications are not completely understood and no specific treatment strategies are available to this date.

Taking this into account, numerous efforts have aimed to better understand the mechanisms used by the virus during infection in order to identify new therapeutic targets. Specifically, several studies have presented evidence implicating ion channels in HIV infection (Raber et al. 1996; Holden et al. 1999; Gelman et al. 2004; Herman et al. 2010; Chen et al. 2011; Liu et al. 2013; Tewari et al. 2015), suggesting that they could be used as potential pharmacological targets. Among these ion channels, nicotinic acetylcholine receptors (nAChRs) have been suggested to play important roles (Bracci et al. 1992; Giunta et al. 2004; González-Lira et al. 2006; Rock et al. 2008; Ballester et al. 2012; Cao et al. 2013; Gundavarapu et al. 2013; Zhang et al. 2015; Báez-Pagán et al. 2015; Ramos et al. 2015; Delgado-Vélez et al. 2015; Liu et al. 2017; Ekins et al. 2017; Capó-Vélez et al. 2018); however, these have not yet been extensively explored. In

Coral M. Capó-Vélez and Manuel Delgado-Vélez have contributed equally to this work.

✉ José A. Lasalde-Dominicci
jlasalde@gmail.com

¹ Department of Biology, University of Puerto Rico, Río Piedras Campus, PO Box 23360, San Juan, PR 00931, USA

² Molecular Sciences Research Center, San Juan, PR 00926, USA

³ Department of Physical Sciences, University of Puerto Rico, Río Piedras Campus, PO Box 23323, San Juan, PR 00931, USA

particular, the $\alpha 7$ -nAChR is emerging as an important player in the HIV field, since it is expressed not only in the brain, but also in a wide variety of immune cells that are targeted during HIV infection, such as macrophages, monocytes, B-lymphocytes, and T-lymphocytes (CD4⁺) (Wang et al. 2003; van der Zanden et al. 2012; Kawashima et al. 2015), making it a suitable target for treatment development.

Due to the presence of $\alpha 7$ -nAChR on both neuronal and immune cells, and its known participation in HIV settings, this review will focus on the role of $\alpha 7$ -nAChRs in two different but equally important aspects of HIV: inflammation (neuroinflammation and peripheral inflammation) and HAND, both of which severely compromise the quality of life of patients infected with HIV. Here, we review the role of $\alpha 7$ -nAChRs in these HIV-derived consequences and suggest the receptor as a possible pharmacological target to treat/ameliorate both HAND and inflammation.

HIV, cART, and HAND: the Current Status

HIV infection continues to be a major global public health issue, having claimed more than 35 million lives so far (UNAIDS 2017). The currently available treatment, i.e., cART, has substantially decreased AIDS-related hospital admissions (Center for Disease Control and Prevention 2014) and changed the course of HIV infection from a defining disease to a manageable condition. Consequently, treating patients with cART declined AIDS-related deaths, decreased maternal-infant viral transmission, and decreased the incidence of HIV-associated dementia (HAD) (Maschke et al. 2000; Saylor et al. 2016).

Although cART is greatly beneficial (INSIGHT START Study Group et al. 2015), it has somewhat limited therapeutic results including long-term toxicity, dependence on patient's daily adherence, and drug–drug interactions, among others (Solomon and Sax 2015). In fact, despite the decrease in viral load and improved or normal CD4⁺ T-cell counts (Antinori et al. 2007), patients still suffer from chronic inflammation (Appay and Sauce 2008; Funderburg et al. 2014); thus, the need for new and innovative therapies is clearly of the utmost importance.

Inflammation is a defense mechanism against diverse insults, and it is programmed to eliminate or remove exogenous noxious agents. This process is mediated by effector molecules such as cytokines, which are released by immune cells during immune responses. Cytokines are key to innate immune responses; however, excessive cytokine production can give rise to chronic or uncontrolled inflammatory responses, an event related to the pathology of many inflammatory diseases (Johnston and Webster 2009; Wang et al. 2009). Inflammation is controlled through the cholinergic anti-inflammatory pathway (CAP), which

seems to be altered in HIV patients (Delgado-Vélez et al. 2015). In non-infected patients, impairment or disruption of the CAP is associated with delirium, depression, ulcerative colitis, postoperative cognitive decline, inflammatory bowel disease, hemorrhagic shock, pancreatitis, exacerbation of ventilator-induced lung injury, septic peritonitis, hypotensive shock, and cardiopulmonary arrest (Lindgren et al. 1993; Borovikova et al. 2000; van Westerloo et al. 2005, 2006; Das 2007; Tracey 2009; dos Santos et al. 2011; Norman et al. 2011, Su et al. 2012; Cerejeira et al. 2012).

Furthermore, around 15–50% of infected individuals develop some form of cognitive and/or motor dysfunctions, collectively known as HAND (Heaton et al. 2010, 2011; Harezlak et al. 2011; Spudich 2013; McGuire et al. 2014; Saylor et al. 2016). It is a frequent complication of HIV infection that causes several neurological disorders including dementia, myelopathy, peripheral neuropathy, and myopathy (Price and Perry 1994; Lindl et al. 2010). HAND is a heterogeneous complication of HIV, being HAD the most severe form of neurological alteration. At one time, HAD was estimated to affect 20–30% of infected patients, but after therapies became available and cART was introduced, this percentage decreased significantly (Heaton et al. 2011). cART treatment predominantly shifted HAND neurological manifestations to milder forms of cognitive impairment instead of the frank dementia seen in the pre-cART era (Antinori et al. 2007). Although antiretroviral therapy has been helpful in the treatment of HAD, there is a therapeutic gap within the milder form of the disease, where cART is less effective (McGuire et al. 2014). The milder forms of HAND, minor cognitive dysfunction (MCD), and asymptomatic neurocognitive impairments (ANI), are estimated to affect around 15–50% of HIV⁺ patients (Simioni et al. 2010; Heaton et al. 2010; Harezlak et al. 2011; Tavazzi et al. 2014; Saylor et al. 2016). This highlights the need to develop innovative pharmacological strategies because therapies aimed to decrease or abrogate HAND cannot necessarily depend on drugs that decrease HIV replication.

The brain is one of the organs most affected by HIV infection (Woods et al. 2009). This reflects the fact that several cells present in the brain, mainly microglia, can be actively infected by HIV since they express receptors needed for infection (Price and Perry 1994). HIV does not infect neurons; however, HAND patients experience a remarkable loss of neurons due to interaction with viral particles, which trigger the release of inflammatory cytokines (Kaul et al. 2001). Neuronal injury is believed to result from a direct effect and/or interaction between viral proteins and the cell, and indirect effects, which are mediated by activated macrophages that secrete neurotoxic factors such as TNF- α , IL-6, and IL-1 β . These neurotoxic factors and/or cytokines can interact with cell surface receptors in neurons and/or astrocytes

and activate signaling pathways that could lead to apoptosis (Saylor et al. 2016).

Thus, HAND is an important cause of cognitive dysfunction and, together with chronic inflammation, remains an important unresolved issue stemming from HIV. Both are equally important aspects that require urgent attention to counteract their chronic deleterious effects.

Nicotinic Acetylcholine Receptors

Nicotinic acetylcholine receptors are fast ionotropic ion channel receptors that bind to and are activated by their endogenous ligands acetylcholine (ACh) and choline; they also bind nicotine (Gahring and Rogers 2005). Neuronal nAChRs include α ($\alpha 2$ – $\alpha 10$) and β subunits ($\beta 2$ – $\beta 4$) arranged in different stoichiometries to form functional receptors (e.g., $\alpha 4\beta 2$, $\alpha 2\beta 2$, $\alpha 2\beta 4$, $\alpha 3\beta 2$, and $\alpha 4\beta 4$, among others). These heteromeric receptors (assembled with more than one type of subunit) are characteristic for their high affinity to nicotine but differ in their pharmacological properties depending on subunit composition (Gahring and Rogers 2005).

The nAChR subunits $\alpha 2$, $\alpha 3$, $\alpha 4$, $\alpha 10$, $\beta 2$, and $\beta 4$ are not able to form functional ion channels by themselves. Conversely, subunits $\alpha 7$ – $\alpha 9$ form homomeric receptors (assembled with only one type of subunit) and comprise a family of receptors that bind nicotine with lower affinity than heteromeric receptors but bind the antagonist snake toxin, α -bungarotoxin (α -bgtx), with high affinity (Breese et al. 1997; Rangwala et al. 1997). Moreover, studies have shown that nAChRs can modulate cellular processes in the central nervous system (CNS). For instance, nicotine administration can improve cognitive performance in normal subjects and produce a number of physiological effects (Ochoa and Lasalde-Dominicci 2007).

Studies performed by Bracci et al. demonstrated a significant homology between a specific sequence of gp120, the coat protein of HIV, and the putative active sites of snake curare-mimetic neurotoxins, which have the ability to bind irreversibly to nAChRs. Furthermore, the authors demonstrated that recombinant gp120 inhibits the binding of the nAChR antagonist, α -bgtx, suggesting that other type of receptors (such as nAChRs) can function as HIV receptors, and supports the notion that ion channels may have a role during HIV infection. Even though this study suggested a potential role for nAChRs in HIV, no specific nAChR subtype was implicated (Bracci et al. 1992). Subsequent studies performed in SIV-infected monkeys found that reduced cholinergic neurotransmission was present in the form of a dramatic reduction in choline acetyltransferase activity, the enzyme responsible for the biosynthesis of ACh (Koutsilieris et al. 2000). Consistent with these observations,

González-Lira et al. (2006) performed electrophysiological studies in rats and found that, in addition to motor and cognition impairments, gp120 interferes with cholinergic neurotransmission as part of the neuropsychiatric abnormalities characteristic of HAD. However, nicotine administration maintained these parameters near normality, supporting the participation of nAChRs in the beneficial outcome. An interesting proof-of-concept study was designed and carried out to evaluate if an increase in the bioavailability of ACh, through the chemical inhibition of acetylcholinesterase, was capable of enhancing the anti-inflammatory reflex in patients infected with HIV (Valdés-Ferrer et al. 2009). Results demonstrate that pyridostigmine (an acetylcholinesterase inhibitor) was able to modify the overactivation and proliferation of T-lymphocytes in patients chronically infected with HIV. This approach also led to an increase in the anti-inflammatory cytokine IL-10, a decrease in T-lymphocytes proliferation, and the production of the proinflammatory cytokine IFN- γ . Moreover, recently, a 16-week proof-of-concept open trial was performed using pyridostigmine as add-on therapy in seven HIV-infected patients with discordant immune response receiving combined antiretroviral therapy, to determine whether pyridostigmine would promote an increase in total CD4⁺ T-cells (Valdés-Ferrer et al. 2017). Results indicate that in patients suffering from HIV⁺, add-on pyridostigmine results in a significant and persistent increase in circulating CD4⁺ T-cells.

Altogether, these studies suggested a role for nAChRs in HIV. However, considering the pharmacological properties and different roles of the $\alpha 7$ -nAChR, a closer look is justified. Below we describe the proposed functions/roles of this receptor to then highlight studies specifically relating to HIV-induced inflammation and HAND, where $\alpha 7$ -nAChRs play a critical part.

The $\alpha 7$ -nAChR: A Unique Ion Channel with Multiple Roles

Many studies have identified roles in health and disease for the $\alpha 7$ -nAChR (Ochoa and Lasalde-Dominicci 2007; Counts et al. 2007; Duffy et al. 2009). The $\alpha 7$ -nAChR is a ligand-gated ion channel that is rapidly desensitized in the presence of a ligand (Conejero-Goldberg et al. 2008). This neuronal receptor is found predominantly in the cerebral cortex, thalamus, hippocampus, basal ganglia, and hypothalamus (both pre- and post-synaptically) (Vázquez-Palacios and Bonilla-Jaime 2004), and immune cells (Wang et al. 2003; Yoshikawa et al. 2006; van der Zanden et al. 2012; Kawashima et al. 2015). Furthermore, $\alpha 7$ -nAChRs can be localized in the soma, presynaptic terminals (regulating neurotransmitter release), and postsynaptic terminals (modulating neuronal excitability) (Gahring and Rogers 2005). Once activated,

its ion pore opens to permit the flow of Na^+ , K^+ , and particularly Ca^{2+} .

The mechanisms by which the $\alpha 7$ -nAChR modulates neurotransmitter release and neuronal excitability are distinct from those proposed for other nAChRs and reflect the channel's unique pharmacological properties. For example, the $\alpha 7$ -nAChR binds α -bgtx with high affinity ($K_d = 94$ pM) (Rangwala et al. 1997) and has low affinity for agonists such as nicotine (Mansvelder et al. 2009). Furthermore, at low agonist concentrations, activation of $\alpha 7$ -nAChR induces a Ca^{2+} influx, leading to a Ca^{2+} -induced Ca^{2+} release from intracellular stores (Mansvelder et al. 2009).

In general, the proposed function for $\alpha 7$ -nAChRs in the brain is to modify the excitability of neurons (Paterson and Nordberg 2000). For example, $\alpha 7$ -nAChRs are expressed in the striatum, where they play a role in the function of local neuronal circuits (Martin 2003). A study conducted by Wu et al. (2004) showed that $\alpha 7$ -nAChRs are expressed on neurons from the substantia nigra and the ventral tegmental area and demonstrated that, once activated, $\alpha 7$ -nAChRs can modulate dopamine release. These results suggest an involvement of the $\alpha 7$ -nAChR in dopaminergic neuronal function and drug reinforcement mechanisms, particularly nicotine dependence. Moreover, nicotine stimulates dopamine release in the nucleus accumbens; an effect thought to be mediated by an increase of glutamate release by $\alpha 7$ -nAChRs located on striatal glutamatergic neurons. This stimulates the release of dopamine in the nucleus accumbens (Kaiser and Wonnacott 2000), further supporting the proposed role of $\alpha 7$ -nAChRs as modulators of neurotransmitter release in the CNS. Therefore, $\alpha 7$ -nAChRs can modulate various signaling pathways that converge to fine-tune neuronal circuits and contribute to subtle but crucial aspects of cognitive behavior (Mansvelder et al. 2009). Consequently, the expression of small numbers of $\alpha 7$ -nAChRs at regulatory sites may lead to multiple outcomes in the performance of normal cells, susceptibility to exogenous agents, or participation in processes ranging from neurodegeneration to inflammation (Gahring and Rogers 2005).

In addition to its central role in the CNS, the $\alpha 7$ -nAChR is also present in immune cells. As a result, several efforts have been made to elucidate its role in these cells. It is now clear that the $\alpha 7$ -nAChR is an essential regulator of inflammation in macrophages through the CAP (Tracey 2007). The anti-inflammatory activity of the $\alpha 7$ -nAChR occurs by the ACh-induced activation of the receptor to inhibit the production of proinflammatory cytokines in macrophages, thereby modulating immune responses and the progression of inflammatory diseases, avoiding organ and systemic damage (Tracey 2009). Altogether, these studies highlight the importance of this neuroimmune tract for the well-being of healthy people and accents the $\alpha 7$ -nAChR's pivotal role in the innate immune response necessary to resolve

inflammation. Despite its well-known role in the CAP, the lack of functional (electrophysiological) evidence has raised the question of whether this receptor acts as an ion-conducting channel (Skok 2009). However, we recently demonstrated that, in human monocyte-derived macrophages (MDMs), the $\alpha 7$ -nAChR retains its ion channel translocation activity (Báez-Pagán et al. 2015). Whether translocated ions play a role in the CAP remains largely unknown (Fig. 1).

Even though the anti-inflammatory role of $\alpha 7$ -nAChR in monocytes and macrophages is well recognized (Wang et al. 2003; Yoshikawa et al. 2006), its role in other immune cells is not well understood. Recent reports demonstrate that $\alpha 7$ -nAChR plays an essential role in the immune synapse between human T- and B-lymphocytes since they are recruited to these sites to inhibit CD40-related mitogenic functions (Kalashnyk et al. 2012). It also regulates B-lymphocytes' proliferation (Koval et al. 2009), development, and survival (Skok et al. 2006). Moreover, a recent study identified the $\alpha 7$ -nAChR as a modulator of lymphocyte activation (De Rosa et al. 2009). In the case of regulatory T-lymphocytes (Tregs), the $\alpha 7$ -nAChR seems to be a critical regulator of the immunosuppressive function of CD4^+ CD25^+ Tregs (Wang et al. 2010). However, it is also important to mention that, in addition to $\alpha 7$ -nAChRs, monocytes, macrophages, B-lymphocytes, and T-lymphocytes also express other homomeric nAChRs, including $\alpha 3$, $\alpha 5$, $\alpha 9$, and $\alpha 10$ (Wang et al. 2003; Kawashima et al. 2012), receptors that could contribute to the net cellular response of these cells during health and disease.

As a whole, this demonstrates the importance of $\alpha 7$ -nAChRs in various cellular processes and highlights its versatility. Thus, this receptor is an integral player of innate processes ranging from the brain to immune cells having clear repercussions in neurotransmitter release, neurodegenerative diseases, and inflammation.

HIV-Induced Inflammation/ Neuroinflammation and $\alpha 7$ -nAChR

The general notion is that in patients infected with HIV, the inflammatory response is exaggerated, uncontrolled, and chronic. Undoubtedly, patients infected with HIV suffer from chronic and persistent inflammatory processes, which cause numerous AIDS and non-AIDS-related complications such as neurocognitive deterioration, cardiovascular disease, thromboembolic disease, type II diabetes, cancer, osteoporosis, multiple end-organ disease, and frailty (Deeks et al. 2013; Barré-Sinoussi et al. 2013; Hearps et al. 2014), even when patients adhere to cART regimens that minimize viral replication and increase CD4 cell counts (Appay and Sauce 2008; Funderburg et al. 2014).




Fig. 1 Effects of HIV infection on macrophages' $\alpha 7$ -nAChRs. **a** In uninfected macrophages, SDF-1 α promotes signaling through CXCR4, a G-protein coupled receptor. In addition, as mentioned before, $\alpha 7$ -nAChR is sensitive to the antagonist α -bgtx. A robust body of evidence demonstrates that the net effect of $\alpha 7$ -nAChR activation is inhibition of the production of cytokines (Wang et al. 2003), consequently decreasing inflammation (efficient CAP). Whether the ion influx through $\alpha 7$ -nAChR is relevant for the proper operation of the CAP or other important physiological processes remains obscure. **b** In infected macrophages, HIV-infected CD4⁺ T-lymphocytes secrete virions and gp120_{IIIb} that interact and stimulate CXCR4 to induce the upregulation of $\alpha 7$ -nAChR (Delgado-Vélez et al. 2015). This upregulation does not result in cell death, or apoptosis, which

is consistent with the anti-apoptotic signature expressed by monocytes recovered from patients infected with HIV (Giri et al. 2009) and those from HIV-infected macrophages (Swingler et al. 2007). Contrary to what was expected, the activation of this high number of $\alpha 7$ -nAChRs with ACh does not enhance the CAP response because their activation does not lead to inhibition of cytokine production, therefore, conferring macrophages a proinflammatory phenotype indicative of a CAP disruption. Moreover, application of an antagonist of $\alpha 7$ -nAChR, bupropion, partially restores the CAP by reducing inflammatory chemokines, but not interleukins (Delgado-Vélez et al. 2015). Interestingly, whether the identity of the non-selective calcium channel previously described (Liu et al. 2000) is $\alpha 7$ -nAChR remains to be determined

In the CNS, the relationship between neuroinflammation and $\alpha 7$ -nAChRs in the context of HIV has been scarcely examined. Available evidence suggests that activation of $\alpha 7$ -nAChRs expressed by microglia (brain macrophages) plays a key role during neuroinflammation processes since its activation seems to suppress inflammation through the CAP, the same mechanism used by macrophages, thus conferring neuroprotective properties (Shytle et al. 2004; Suzuki et al. 2006). Giunta et al. (2004) reported an in vitro model of HIV-associated dementia composed of cultured microglia cells synergistically activated by IFN- γ and gp120. However, this activation was synergistically attenuated through $\alpha 7$ -nAChRs and the

p44/42 MAPK pathway by pretreatment with nicotine and the cholinesterase inhibitor, galantamine.

The detrimental consequences of HIV infection are not limited to the presence of viral constituents in the body, such as gp120. Accumulating evidence also demonstrates that neuroinflammation could emerge from soluble virions, HIV-infected monocytes, and T-lymphocytes that cross the permeabilized blood–brain barrier (BBB). These infected cells not only affect brain-resident cells upon migration into the CNS but also produce proinflammatory cytokines such as TNF- α and IL-1 β , which in turn, activate microglia and astrocytes. Cytokine levels in the cerebrospinal fluid are associated with neurocognitive impairment in

HIV infection. Activated brain-resident cells (e.g., microglia), along with perivascular macrophages, are the main contributors to neuroinflammation in HIV infection (Hong and Banks 2015). They can release neurotoxic factors, such as excitatory amino acids and inflammatory mediators, resulting in neuronal dysfunction and death (Hong and Banks 2015). Moreover, viral proteins (e.g., gp120), which induce brain endothelial cells to release cytokines, may also contribute to brain inflammation and the pathogenesis of HAND. In fact, gp120 facilitates infiltration of HIV-infected immune cells due to permeabilization of the BBB (Kanmogne et al. 2007). Interestingly, a recent study was conducted to evaluate the effects of antiretroviral therapy on the brain. Results showed that indinavir (a protease inhibitor) acts as a positive allosteric modulator of $\alpha 7$ -nAChRs at low concentrations, whereas at high concentrations, it acts as an inhibitor of $\alpha 7$ -nAChRs in the brain, thus possibly contributing to the cognitive dysfunction observed in patients infected with HIV (Ekins et al. 2017). However, caution is needed regarding the possible effects of antiretroviral therapy on the CNS.

Another report on this matter demonstrated that the $\alpha 7$ -nAChR agonists, choline, and GTS-21 attenuated the intrathecal gp120-induced proinflammatory cytokine- and microglia-dependent mechanical allodynia (Loram et al. 2010). Of note, choline (full agonist of $\alpha 7$ -nAChR) significantly blocked and reversed the gp120-induced mechanical allodynia for at least 4 h after administration. In addition, intrathecal choline delivered either with or 30 min after gp120 reduced the gp120-induced production of IL-1 β protein and proinflammatory cytokine mRNAs within the lumbar spinal cord. The second $\alpha 7$ -nAChR agonist, GTS-21, also significantly reversed the gp120-induced mechanical allodynia and lumbar spinal cord levels of proinflammatory cytokine mRNAs and IL-1 β protein.

The CAP is an innate neuroimmune mechanism whereby cytokine production is inhibited via activation of $\alpha 7$ -nAChRs expressed in macrophages; this avoids tissue and organ damage due to an exaggerated and persistent inflammation (Tracey 2002; Wang et al. 2009). This pathway provides a fast, discrete, and localized way of controlling immune responses and regulating inflammation (Johnston and Webster 2009) triggered by proinflammatory molecules (e.g., cytokines, LPS). Upon its detection, a signal travels to the CNS via the afferent vagus nerve fibers and stimulates the release of ACh by the efferent arm of the vagus nerve. Then, ACh activates $\alpha 7$ -nAChRs in macrophages to inhibit cytokine production and balance the inflammatory response (Tracey 2002, 2009; Wang et al. 2009). In addition, electrical stimulation of the vagus nerve and $\alpha 7$ -nAChR agonists ameliorate symptoms in models of inflammatory diseases, demonstrating a pivotal role for $\alpha 7$ -nAChRs in the regulation of inflammation.

Interestingly, this mechanism is conserved in microglia, where $\alpha 7$ -nAChRs regulate neuroinflammation (Shytle et al. 2004). Because both macrophages and microglia regulate inflammation through the same mechanism, it could be speculated that they could regulate each other. Indeed, a recent study demonstrates that electrical activation of the vagus nerve decreases the levels of proinflammatory cytokines and the percent of activated microglia in the brain (Meneses et al. 2016). These results provide an important piece of information since neuroinflammation is a hallmark of several infections and neurodegenerative diseases. Thus, strategies to control neuroinflammation could result in a definite improvement in patients (Meneses et al. 2016).

Although the importance of the CAP is evident, so far it has not been studied extensively in patients infected with HIV. Recently, our group examined the cholinergic anti-inflammatory response (CAR) in HIV. We found that gp120_{IIIIB} addition to MDMs induces upregulation of $\alpha 7$ -nAChRs in vitro (Fig. 1). Moreover, inflammation studies performed in MDMs pre-exposed to gp120_{IIIIB} and challenged with LPS demonstrated that ACh addition is no longer able to reduce inflammation, suggesting a CAR disruption. Furthermore, examination of $\alpha 7$ -nAChR levels in monocytes, MDMs, and T-lymphocytes recovered from patients infected with HIV revealed a remarkable upregulation in these cell types (Delgado-Vélez et al. 2015). Finally, application of bupropion (an $\alpha 7$ -nAChR antagonist) to MDMs upregulated for $\alpha 7$ -nAChR (by gp120) partially restores the CAR by reducing inflammatory chemokines, but not interleukins (Delgado-Vélez et al. 2015) (Fig. 1).

Overall, HIV-induced inflammation occurs in two scenarios: the periphery and the CNS. Nevertheless, the specific mechanisms by which they occur are not entirely understood, and further studies are needed to identify new therapeutic targets to generate new neuroprotective medications. The studies described here highlight the importance of considering the $\alpha 7$ -nAChR and the CAP in HIV-induced pathologies and expand the current knowledge about neuroimmunomodulation processes during HIV infection.

HAND and the $\alpha 7$ -nAChR

The $\alpha 7$ -nAChR is amply distributed through the brain. In a series of works, the $\alpha 7$ -nAChR has been implicated in different HIV-related settings. It was shown that gp120 injected into the hippocampus impairs memory by affecting the cholinergic/vasoactive intestinal peptide system (Farr et al. 2002). More recently, a report implicated $\alpha 7$ -nAChRs in Cryptococcal-induced meningitis, the most common fungal opportunistic infection in the CNS of patients infected with HIV (Zhang et al. 2015). Specifically, the authors demonstrated that $\alpha 7$ -nAChRs play a detrimental role in the host's

defense against *Cryptococcus neoformans* and HIV-1 associated comorbidity factors that compromise BBB integrity (Zhang et al. 2015). Furthermore, a recent report suggests that $\alpha 7$ -nAChR may play an important role in the neuropathology caused by gp120, methamphetamine, and nicotine, which are the major pathogenic factors contributing to the pathogenesis of HAND (Liu et al. 2017). Moreover, Gundavarapu et al. (2013) demonstrated that HIV-gp120 induces and regulates mucus formation on airway epithelial cells through a CXCR4- $\alpha 7$ -nAChR-GABA_AR-dependent pathway.

In Ballester et al. (2012), a potential role for $\alpha 7$ -nAChR in HAND was established, as the HIV-1 coat protein gp120_{IIIIB} was shown to induce a functional upregulation of $\alpha 7$ -nAChRs promoting an increase in Ca²⁺ entry that resulted in neuronal cell death (Fig. 2). Along this line, Ramos et al. (2015) showed that, in neuronal cells, gp120_{IIIIB} is able to increase the expression of the $\alpha 7$ -nAChR gene, *CHRNA7*, together with a down-regulation of *CHRFAM7A*, the partial duplication of the $\alpha 7$ -nAChR gene (dup $\alpha 7$) (Fig. 2). Moreover, evaluation of these genes in the basal ganglia of patients infected with HIV suffering from different stages of HAND

demonstrates an alteration in the *CHRFAM7A/CHRNA7* expression ratio. Finally, Capó-Vélez et al. (2018) demonstrated that, in mice expressing gp120_{IIIIB} in the brain, striatal neurons are upregulated for $\alpha 7$ -nAChRs, and its activation leads to an increase in intracellular calcium and eventual apoptosis, an effect that can be attenuated by antagonizing the $\alpha 7$ -nAChR (Fig. 2). Moreover, they showed that gp120_{IIIIB} mice perform worse than wild-type mice on a striatum-dependent behavioral task and demonstrate locomotor impairments, which improve significantly with $\alpha 7$ -nAChR antagonists (Capó-Vélez et al. 2018).

Taken together, the available literature (summarized in Table 1) highlights the importance of studying the role of the $\alpha 7$ -nAChR in an HIV-related setting and demonstrates its feasibility as a pharmacological target.

Concluding Remarks

HIV infection represents a major burden to the immune system. Decades of basic and clinical research have made it clear that HIV infection promotes uncontrolled inflammation in both the periphery and the CNS of infected




Fig. 2 Effects of HIV infection on neurons' $\alpha 7$ -nAChRs. 1–4 gp120 released from HIV-infected microglia interact with CXCR4 expressed by neurons, activating the Ras/Raf/MEK/Erk/Egr-1 signaling pathway that eventually promotes $\alpha 7$ -nAChR upregulation in neuronal cells (Ballester et al. 2012). Moreover, this response can be recapitulated by adding SDF-1 α , the endogenous agonist of CXCR4, which also induces $\alpha 7$ -nAChR overexpression in neuronal cells (Ballester

et al. 2012). These responses are sensitive to CXCR4 blockade with AMD3100 and MEK inhibition by U0126 or PD98059. 5 Furthermore, $\alpha 7$ -nAChR activation with ACh promotes intracellular Ca²⁺ overload that results in cell death and apoptosis (Ballester et al. 2012; Capó-Vélez et al. 2018). 6 The gp120-induced $\alpha 7$ -nAChR upregulation described in points 1–5 involves a decrease in dup $\alpha 7$ expression, its repressive partial duplication (Ramos et al. 2015)

Table 1 Studies that establish relationships between HIV and $\alpha 7$ -nAChR

nAChR or subtype	Finding(s)	Relevance to HIV field	Reference
nAChRs	A significant homology between HIV-gp120 and snake curare-mimetic neurotoxins, which specifically bind to nAChRs was found	The first suggestion was that HIV-1 gp120 could bind to nAChR expressed in muscle cells and neurons. In addition, the nAChR was proposed as an HIV receptor	
	Nicotine, a nAChR agonist, enhances production of HIV of in vitro-infected alveolar macrophages from healthy cigarette smokers	HIV-infected smokers should experience an increase in viral load	Farr et al. (2002)
	HIV-gp120 injected in the hippocampus impairs memory in rats, an effect reversed by hippocampal cholinergic stimulation	Cholinergic pathways could have an essential role in reversing the cognitive decline induced by gp120 neurotoxicity	González-Lira et al. (2006)
	The coat protein gp120 is involved in the pathogenesis of neuropsychiatric abnormalities; it occurs by interfering with cholinergic neurotransmission	Propose a mechanism of action for gp120 by interfering with cholinergic pathways, a phenomenon that could be part of the etiology of cognitive disturbances. Importantly, nicotine treatment seems to be beneficial	
	Nicotine-treated microglia show increased HIV-1 expression in a concentration-dependent manner	nAChRs are important in the neuropathogenesis of CNS in patients infected with HIV	Rock et al. 92008)
	Nicotine treatment restored gp120-impaired pathways in the brain of HIV-transgenic rats including the prefrontal cortex, dorsal hippocampus, and dorsal striatum	Nicotine has beneficial effects on HIV-induced neurological deficits, presumably through the activation of nAChRs in the brain	Cao et al. (2013)
$\alpha 7$ -nAChRs	$\alpha 7$ -nAChR is expressed in macrophages, where it is an essential regulator of inflammation acting through the cholinergic anti-inflammatory pathway	Since chronic inflammation is a common phenomenon in patients infected with HIV (despite good viral control and CD4 ⁺ recovery), the $\alpha 7$ -nAChR could be an important target for the regulation of inflammation	Wang et al. (2003)
	In cultured microglia activated with both interferon and gp120 (an in vitro model of HAD), microglial activation was attenuated by pretreatment with nicotine and galantamine. This occurred through an $\alpha 7$ -nAChR and p44/42 MAPK-dependent mechanism	Novel therapeutic combination to treat or prevent HAD development through modulation of microglial activation using $\alpha 7$ -nAChR agonists and/or acetylcholinesterase inhibitors	Giunta et al. (2004)
	Choline, an $\alpha 7$ -nAChR-specific agonist (Alkondon et al. 1997), significantly blocked and reversed gp120-induced mechanical allodynia	$\alpha 7$ -nAChR as a potential target to modulate and/or abrogate gp120-induced allodynia	Loram et al. (2010)
	Exposure of neuronal cells to gp120 induces a functional $\alpha 7$ -nAChR upregulation that leads to an increase in calcium influx and subsequent cell death. The $\alpha 7$ -nAChR upregulation was also observed in a gp120-transgenic mouse model, specifically in the striatum (basal ganglia), a region greatly affected in patients infected with HIV	Suggest a novel pathway for gp120-induced neuronal cell death. Results position the $\alpha 7$ -nAChR as a potential therapeutic target to ameliorate or reverse HAND-related cognitive dysfunction since antagonists for this receptor are commercially available	Ballester et al. (2012)
	Normal human broncoepithelial cells produce mucus in response to CXCR4-tropic gp120. Mucus formation was blocked by CXCR4, $\alpha 7$ -nAChR, and GABA _A R inhibitors	First demonstration that HIV-gp120 induces and regulates mucus formation in the airway epithelial cells through the CXCR4- $\alpha 7$ -GABA _A R pathway, which may contribute to the higher incidence of obstructive pulmonary diseases in this population	Gundavarapu et al. (2013)

Table 1 (continued)

nAChR or subtype	Finding(s)	Relevance to HIV field	Reference
	In neuronal cells where the $\alpha 7$ -nAChR is upregulated after gp120 exposure, measurements of $\alpha 7$ -nAChR expression gene (<i>CHRNA7</i>) and its partial duplication, dup $\alpha 7$, (<i>CHRFAM7A</i>) reveal that the <i>CHRNA7</i> increase is accompanied by a reduction of its repressive partial duplication, <i>CHRFAM7A</i> . Moreover, there is a dysregulation of evaluation of <i>CHRNA7</i> and <i>CHRFAM7A</i> expression in postmortem brain samples from patients infected with HIV suffering from different stages of HAD	Shown dysregulation in <i>CHRFAM7A/CHRNA7</i> expression ratio in the basal ganglia from postmortem brain samples of HIV-infected subjects. These results coupled with the high calcium permeability of $\alpha 7$ -nAChR may explain the significant loss of neurons in this area, which is a common manifestation of HIV-induced neurological disorders (Everall et al. 1995; Brew et al. 1995)	Ramos et al. (2015)
	Both chemical and genetic blockage of $\alpha 7$ -nAChRs are protective against <i>C. neoformans</i> - and HIV-1 associated comorbidity factors-induced BBB injury and CNS disorders by down-regulation of circulating brain microvascular endothelial cells (cBMEC) shedding, monocyte recruitment, NF- κ B signaling, senescence, and neuronal inflammation	Pharmacological blockade of $\alpha 7$ -nAChR in the BBB could prevent HIV-1 associated comorbidity factors-induced BBB injury and CNS disorders	Zhang et al. (2015)
	The HIV Env gp120 induces the upregulation of $\alpha 7$ -nAChR in MDMs. In addition, patients infected with HIV are upregulated for $\alpha 7$ -nAChR in their monocytes, MDMs, and T-lymphocytes. Activation of the cholinergic anti-inflammatory response in $\alpha 7$ -nAChR-upregulated MDMs shows that gp120 disrupts this innate immune mechanism to control inflammation. Furthermore, application of an $\alpha 7$ -nAChR antagonist, bupropion, partially restores the response	$\alpha 7$ -nAChR emerges as a potential pharmacological target to control chronic inflammation in HIV-infected subjects	Delgado-Vélez et al. (2015)
	They demonstrate that blockade of $\alpha 7$ -nAChR could significantly reduce HIV-1 gp120, methamphetamine, and nicotine-induced BBB injury and CNS disorders by decreasing amyloid-beta transport, leukocyte recruitment, cholinergic signaling, premature senescence of brain microvascular endothelial cells, and neuroinflammation	This work suggests that $\alpha 7$ -nAChR may play an essential role in neuropathology caused by gp120, methamphetamine, and nicotine, which are the major pathogenic factors contributing to the pathogenesis of HAND	Liu et al. (2017)
	This group demonstrated that at low concentrations, indinavir (a protease inhibitor), acts as a positive allosteric modulator of $\alpha 7$ -nAChR, whereas at concentrations greater than 10 mmol/l indinavir acts as an inhibitor of the $\alpha 7$ -nAChR	This work demonstrates that an antiretroviral therapeutic drug interferes with the $\alpha 7$ -nAChR in the brain, thus possibly contributing to the cognitive dysfunction observed in patients infected with HIV. Moreover, it opens a venue to investigate the effects of other antiretrovirals on $\alpha 7$ -nAChR and expand the efforts to other cholinergic receptors involved in cognition	Ekims et al. (2017)
	In a murine model of gp120 _{11B} expressed in the brain, it was found that $\alpha 7$ -nAChR is overexpressed in striatal neurons and its activation promotes apoptosis. Moreover, a striatum-dependent task showed that these animals have a poorer performance and exhibit locomotor impairment. Furthermore, administration of $\alpha 7$ -nAChR antagonists improved the locomotor performance in these animals significantly	Consistent with Zhang et al. (2015) and Delgado-Vélez et al. (2015), results of this work position $\alpha 7$ -nAChR as an attractive pharmacological target that should be exploited in the HIV field	Capó-Vélez et al. (2018)

subjects. Furthermore, HIV infection does not only compromise the patient's cognitive functions (as part of HAND development) but also affects peripheral organs as a result of chronic and persistent inflammation. Although significant advances have been made regarding HIV-induced pathologies, an effective adjunctive therapy to mitigate chronic inflammation to avoid damage to organs and/or HAND development remains elusive. The inflammation suffered by patients with HIV represents a major clinical challenge because it is necessary to rectify disproportioned inflammation processes in the CNS and periphery simultaneously. An anti-inflammatory medication that crosses the BBB and acts in the periphery could be an ideal strategy to decrease inflammation on the CNS and periphery in patients infected with HIV. Moreover, a pharmacological target expressed in both CNS and periphery represents a unique opportunity to counteract neuroinflammation/peripheral inflammation and to prevent HAND as well as the appearance of non-AIDS-related diseases. Indeed, this is what the $\alpha 7$ -nAChR offers: a new therapeutic target for adjunctive therapy expressed in the brain and immune cells. In the brain, $\alpha 7$ -nAChR activation improves cognitive processes (Thomsen et al. 2010) and decreases neuroinflammation (Egea et al. 2015), whereas in the periphery, $\alpha 7$ -nAChR activation in macrophages and monocytes inhibits the production of proinflammatory cytokines (Wang et al. 2003; Yoshikawa et al. 2006). The inflammatory process that is developed in the brain of patients infected with HIV has been the focus of recent studies. Neuroinflammation occurs when microglia, astrocytes, and perivascular macrophages are activated. Therefore, $\alpha 7$ -nAChR clearly represents a unique pharmacological target that could be exploited in the HIV field. Still, it is important to consider that $\alpha 7$ -nAChR is also expressed in other immune cells such as lymphocytes and other non-immune cells in which the net effect of its systemic activation, modulation, or blockade is still unknown.

Because the $\alpha 7$ -nAChR has received much attention by the pharmaceutical industry, many molecules that target this receptor have recently been developed. According to clinicaltrials.gov (consulted June 18, 2017), eleven trials with different drugs targeting $\alpha 7$ -nAChR have been recently executed, including nine completed, one terminated, and one active (recruiting). In addition to the synthesis of new therapeutics, it is also possible to take advantage of FDA-approved drugs available in the market. For instance, the antidepressant and smoking cessation medication bupropion is an FDA-approved $\alpha 7$ -nAChR antagonist that has been used in patients infected with HIV (Park-Wyllie and Antoniou 2003; Currier et al. 2003; Pedrol-Clotet et al. 2006). Importantly, in HIV-negative individuals and animal models, bupropion use has demonstrated anti-inflammatory properties (Altschuler and

Kast 2003; Brustolim et al. 2006). Furthermore, bupropion has been shown to decrease some chemokines, but not interleukins, in human macrophages upregulated for $\alpha 7$ -nAChR by the HIV-gp120_{IIIB} (Delgado-Vélez et al. 2015) (Fig. 1). Indeed, drugs that agonize the $\alpha 7$ -nAChR with anti-inflammatory benefits have been described (de Jonge and Ulloa 2007; Pohanka 2012), and new potential compounds continuously emerge (Sahdeo et al. 2014; van Maanen et al. 2015).

Another plausible medication is varenicline, a full agonist of $\alpha 7$ -nAChRs (Mihalak et al. 2006) and smoking cessation drug that has been demonstrated to be safe and effective in patients infected with HIV (Cui et al. 2012; Ferketich et al. 2013). Results from varenicline suggest similar or even higher cessation rates than for the general population (Cui et al. 2012; Ferketich et al. 2013; Shirley et al. 2013). Thus, repurposing FDA-approved pharmacotherapeutics targeted at $\alpha 7$ -nAChRs to treat HIV-associated diseases implies that new treatments could reach patients faster.

An understudied perspective on HIV-induced chronic inflammation and HAND is the role of nAChRs. Here, we provide evidence demonstrating that the $\alpha 7$ -nAChR is upregulated in HIV settings in a gp120-dependent manner in both neural and immune cells (Ballester et al. 2012; Ramos et al. 2015; Delgado-Vélez et al. 2015; Capó-Vélez et al. 2018) and that HIV-infected monocytes, macrophages, and T-cells are also upregulated in HIV-infected women (Delgado-Vélez et al. 2015). The observed increase in $\alpha 7$ -nAChR levels is in accordance with the $\alpha 7$ -nAChR upregulation reported under inflammation settings in T-lymphocytes, alveolar macrophages, and neutrophils (Su et al. 2007; Chernyavsky et al. 2009; Nizri et al. 2009). Apparently, $\alpha 7$ -nAChR upregulation is a typical response of immune cells during inflammation challenges, whereas in the CNS it appears to be a response induced by the HIV viral glycoprotein, gp120 (Ballester et al. 2012; Capó-Vélez et al. 2018) or the viral presence itself. Overall, the $\alpha 7$ -nAChR offers the opportunity to explore new pharmacological treatments through the development of new drugs or by testing available drugs for HIV treatment.

It is not clear whether high levels of $\alpha 7$ -nAChRs in immune or neuronal cells (particularly microglia and macrophages) represent an anti-inflammatory ally that can be exploited as a therapeutic target since our recent results demonstrate that upregulation of $\alpha 7$ -nAChRs does not potentiate the anti-inflammatory response in macrophages (Delgado-Vélez et al. 2015). Moreover, whether the upregulation observed in neural cells and neural tissues (Ballester et al. 2012; Ramos et al. 2015; Capó-Vélez et al. 2018) represents a pharmacological intervention opportunity remains undetermined. However, positive allosteric modulators of $\alpha 7$ -nAChRs, such as PNU-120596, can be tested to try to rescue $\alpha 7$ -nAChR's anti-inflammatory activity (Hurst

et al. 2005). Thus, efforts to better understand the molecular mechanism that impede the appropriate function of the cholinergic anti-inflammatory response in macrophages in HIV settings are definitely critical in the development of new strategies to reduce neuroinflammation and peripheral inflammation in patients infected with HIV. Moreover, more research is necessary to better understand the role of $\alpha 7$ -nAChR in the neurocognitive deterioration experienced by patients infected with HIV.

Acknowledgements This work was supported by National Institutes of Health-NINDS (Grant No. 2U54NS43011) and COBRE (Grant No. 1P20GM103642) to J.A.L-D, RISE Program (Grant No. 2R25GM061151) to M.D-V and C.M.C-V. The project described was supported by Award Number 2U54MD007587 from the National Institute on Minority Health and Health Disparities and Award Number P30MH075673 from the National Institute of Mental Health to C.A.B-P. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions Both C.M.C-V. and M.D-V. performed literature research and wrote the review. J.A.L-D. and C.A.B-P. helped with the conceptualization of the review and performed literature research.

Compliance with Ethical Standards

Conflict of interest The authors declare that there is no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alkondon M, Pereira EF, Cortes WS et al (1997) Choline is a selective agonist of $\alpha 7$ nicotinic acetylcholine receptors in the rat brain neurons. *Eur J Neurosci* 9:2734–2742
- Althoff KN, Smit M, Reiss P, Justice AC (2016) HIV and ageing: improving quantity and quality of life. *Curr Opin HIV AIDS* 11:527–536. <https://doi.org/10.1097/COH.0000000000000305>
- Altschuler EL, Kast RE (2003) Bupropion in psoriasis and atopic dermatitis: decreased tumor necrosis factor- α ? *Psychosom Med* 65:719
- Antinori A, Arendt G, Becker JT et al (2007) Updated research nosology for HIV-associated neurocognitive disorders. *Neurology* 69:1789–1799. <https://doi.org/10.1212/01.WNL.0000287431.88658.8b>
- Appay V, Sauce D (2008) Immune activation and inflammation in HIV-1 infection: causes and consequences. *J Pathol* 214:231–241. <https://doi.org/10.1002/path.2276>
- Báez-Pagán CA, Delgado-Vélez M, Lasalde-Dominicci JA (2015) Activation of the macrophage $\alpha 7$ nicotinic acetylcholine receptor and control of inflammation. *J Neuroimmune Pharmacol* 10:468–476. <https://doi.org/10.1007/s11481-015-9601-5>
- Ballester LY, Capó-Vélez CM, García-Beltrán WF et al (2012) Up-regulation of the neuronal nicotinic receptor $\alpha 7$ by HIV glycoprotein 120: potential implications for HIV-associated neurocognitive disorder. *J Biol Chem* 287:3079–3086. <https://doi.org/10.1074/jbc.M111.262543>
- Barré-Sinoussi F, Ross AL, Delfraissy J-F (2013) Past, present and future: 30 years of HIV research. *Nat Rev Microbiol* 11:877–883. <https://doi.org/10.1038/nrmicro3132>
- Borovikova LV, Ivanova S, Zhang M et al (2000) Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405:458–462. <https://doi.org/10.1038/35013070>
- Bracci L, Lozzi L, Rustici M, Neri P (1992) Binding of HIV-1 gp120 to the nicotinic receptor. *FEBS Lett* 311:115–118
- Breese CR, Adams C, Logel J et al (1997) Comparison of the regional expression of nicotinic acetylcholine receptor $\alpha 7$ mRNA and [125 I]- α -bungarotoxin binding in human post-mortem brain. *J Comp Neurol* 387:385–398
- Brew BJ, Rosenblum M, Cronin K, Price RW (1995) AIDS dementia complex and HIV-1 brain infection: clinical-virological correlations. *Ann Neurol* 38:563–570. <https://doi.org/10.1002/ana.410380404>
- Brustolim D, Ribeiro-dos-Santos R, Kast RE et al (2006) A new chapter opens in anti-inflammatory treatments: the antidepressant bupropion lowers production of tumor necrosis factor- α and interferon- γ in mice. *Int Immunopharmacol* 6:903–907. <https://doi.org/10.1016/j.intimp.2005.12.007>
- Cao J, Wang S, Wang J et al (2013) RNA deep sequencing analysis reveals that nicotine restores impaired gene expression by viral proteins in the brains of HIV-1 transgenic rats. *PLoS ONE* 8:e68517. <https://doi.org/10.1371/journal.pone.0068517>
- Capó-Vélez CM, Morales-Vargas B, García-González A et al (2018) The $\alpha 7$ -nicotinic receptor contributes to gp120-induced neurotoxicity: implications in HIV-associated neurocognitive disorders. *Sci Rep* 8:1829. <https://doi.org/10.1038/s41598-018-20271-x>
- Center for Disease Control and Prevention (2014) HIV surveillance report 2012
- Cerejeira J, Nogueira V, Luís P et al (2012) The cholinergic system and inflammation: common pathways in delirium pathophysiology. *J Am Geriatr Soc* 60:669–675. <https://doi.org/10.1111/j.1532-5415.2011.03883.x>
- Chen L, Liu J, Xu C et al (2011) HIV-1gp120 induces neuronal apoptosis through enhancement of 4-aminopyridine-sensitive outward K^+ currents. *PLoS ONE* 6:e25994. <https://doi.org/10.1371/journal.pone.0025994>
- Chernyavsky AI, Arredondo J, Galitovskiy V et al (2009) Structure and function of the nicotinic arm of acetylcholine regulatory axis in human leukemic T cells. *Int J Immunopathol Pharmacol* 22:461–472
- Conejero-Goldberg C, Davies P, Ulloa L (2008) $\alpha 7$ nicotinic acetylcholine receptor: a link between inflammation and neurodegeneration. *Neurosci Biobehav Rev* 32:693–706. <https://doi.org/10.1016/j.neubiorev.2007.10.007>
- Counts SE, He B, Che S et al (2007) $\alpha 7$ nicotinic receptor up-regulation in cholinergic basal forebrain neurons in Alzheimer disease. *Arch Neurol* 64:1771–1776. <https://doi.org/10.1001/archneur.64.12.1771>
- Cui Q, Robinson L, Elston D et al (2012) Safety and tolerability of varenicline tartrate (Champix®)/Chantix® for smoking cessation in HIV-infected subjects: a pilot open-label study. *AIDS Patient Care STDs* 26:12–19. <https://doi.org/10.1089/apc.2011.0199>

- Currier MB, Molina G, Kato M (2003) A prospective trial of sustained-release bupropion for depression in HIV-seropositive and AIDS patients. *Psychosomatics* 44:120–125
- Das UN (2007) Vagus nerve stimulation, depression, and inflammation. *Neuropsychopharmacology* 32:2053–2054. <https://doi.org/10.1038/sj.npp.1301286>
- De Jonge WJ, Ulloa L (2007) The alpha7 nicotinic acetylcholine receptor as a pharmacological target for inflammation. *Br J Pharmacol* 151:915–929. <https://doi.org/10.1038/sj.bjp.0707264>
- De Rosa MJ, Dionisio L, Agriello E et al (2009) Alpha 7 nicotinic acetylcholine receptor modulates lymphocyte activation. *Life Sci* 85:444–449. <https://doi.org/10.1016/j.lfs.2009.07.010>
- Deeks SG, Tracy R, Douek DC (2013) Systemic effects of inflammation on health during chronic HIV infection. *Immunity* 39:633–645. <https://doi.org/10.1016/j.immuni.2013.10.001>
- Delgado-Vélez M, Báez-Pagán CA, Gerena Y et al (2015) The $\alpha 7$ -nicotinic receptor is upregulated in immune cells from HIV-seropositive women: consequences to the cholinergic anti-inflammatory response. *Clin Transl Immunol* 4:e53. <https://doi.org/10.1038/cti.2015.31>
- Dos Santos CC dos, Shan Y, Akram A et al (2011) Neuroimmune regulation of ventilator-induced lung injury. *Am J Respir Crit Care Med* 183:471–482. <https://doi.org/10.1164/rccm.201002-0314OC>
- Duffy AM, Zhou P, Milner TA, Pickel VM (2009) Spatial and intracellular relationships between the alpha7 nicotinic acetylcholine receptor and the vesicular acetylcholine transporter in the prefrontal cortex of rat and mouse. *Neuroscience* 161:1091–1103. <https://doi.org/10.1016/j.neuroscience.2009.04.024>
- Egea J, Buendia I, Parada E et al (2015) Anti-inflammatory role of microglial alpha7 nAChRs and its role in neuroprotection. *Biochem Pharmacol* 97:463–472. <https://doi.org/10.1016/j.bcp.2015.07.032>
- Ekins S, Mathews P, Saito EK et al (2017) $\alpha 7$ -Nicotinic acetylcholine receptor inhibition by indinavir: implications for cognitive dysfunction in treated HIV disease. *AIDS Lond Engl* 31:1083–1089. <https://doi.org/10.1097/QAD.0000000000001488>
- Everall I, Barnes H, Spargo E, Lantos P (1995) Assessment of neuronal density in the putamen in human immunodeficiency virus (HIV) infection. Application of stereology and spatial analysis of quadrats. *J Neurovirol* 1:126–129
- Farr SA, Banks WA, Uezu K et al (2002) Mechanisms of HIV type 1-induced cognitive impairment: evidence for hippocampal cholinergic involvement with overstimulation of the VIPergic system by the viral coat protein core. *AIDS Res Hum Retrovir* 18:1189–1195. <https://doi.org/10.1089/08892220260387931>
- Ferketich AK, Diaz P, Browning KK et al (2013) Safety of varenicline among smokers enrolled in the lung HIV study. *Nicotine Tob Res* 15:247–254. <https://doi.org/10.1093/ntr/nts121>
- Funderburg NT, Jiang Y, Debanne SM et al (2014) Rosuvastatin treatment reduces markers of monocyte activation in HIV-infected subjects on antiretroviral therapy. *Clin Infect Dis* 58:588–595. <https://doi.org/10.1093/cid/cit748>
- Gahring LC, Rogers SW (2005) Neuronal nicotinic acetylcholine receptor expression and function on nonneuronal cells. *AAPS J* 7:E885–E894. <https://doi.org/10.1208/aapsj070486>
- Gelman BB, Soukup VM, Schuenke KW et al (2004) Acquired neuronal channelopathies in HIV-associated dementia. *J Neuroimmunol* 157:111–119. <https://doi.org/10.1016/j.jneur.2004.08.044>
- Giri MS, Nebozyhn M, Raymond A et al (2009) Circulating monocytes in HIV-1-infected viremic subjects exhibit an antiapoptosis gene signature and virus- and host-mediated apoptosis resistance. *J Immunol Baltim Md* 1950 182:4459–4470. <https://doi.org/10.4049/jimmunol.0801450>
- Giunta B, Ehrhart J, Townsend K et al (2004) Galantamine and nicotine have a synergistic effect on inhibition of microglial activation induced by HIV-1 gp120. *Brain Res Bull* 64:165–170. <https://doi.org/10.1016/j.brainresbull.2004.06.008>
- González-Lira B, Rueda-Orozco PE, Galicia O et al (2006) Nicotine prevents HIVgp120-caused electrophysiological and motor disturbances in rats. *Neurosci Lett* 394:136–139. <https://doi.org/10.1016/j.neulet.2005.10.021>
- Gundavarapu S, Mishra NC, Singh SP et al (2013) HIV gp120 induces mucus formation in human bronchial epithelial cells through CXCR4/ $\alpha 7$ -nicotinic acetylcholine receptors. *PLoS ONE* 8:e77160. <https://doi.org/10.1371/journal.pone.0077160>
- Harezlak J, Buchthal S, Taylor M et al (2011) Persistence of HIV-associated cognitive impairment, inflammation, and neuronal injury in era of highly active antiretroviral treatment. *AIDS Lond Engl* 25:625–633. <https://doi.org/10.1097/QAD.0b013e3283427da7>
- Hearps AC, Martin GE, Rajasuriar R, Crowe SM (2014) Inflammatory co-morbidities in HIV + individuals: learning lessons from healthy ageing. *Curr HIV/AIDS Rep* 11:20–34. <https://doi.org/10.1007/s11904-013-0190-8>
- Heaton RK, Clifford DB, Franklin DR et al (2010) HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology* 75:2087–2096. <https://doi.org/10.1212/WNL.0b013e318200d727>
- Heaton RK, Franklin DR, Ellis RJ et al (2011) HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *J Neurovirol* 17:3–16. <https://doi.org/10.1007/s13365-010-0006-1>
- Herman BD, Votruba I, Holy A et al (2010) The acyclic 2,4-diaminopyrimidine nucleoside phosphonate acts as a purine mimetic in HIV-1 reverse transcriptase DNA polymerization. *J Biol Chem* 285:12101–12108. <https://doi.org/10.1074/jbc.M109.096529>
- Holden CP, Haughey NJ, Nath A, Geiger JD (1999) Role of Na^+/H^+ exchangers, excitatory amino acid receptors and voltage-operated Ca^{2+} channels in human immunodeficiency virus type 1 gp120-mediated increases in intracellular Ca^{2+} in human neurons and astrocytes. *Neuroscience* 91:1369–1378
- Hong S, Banks WA (2015) Role of the immune system in HIV-associated neuroinflammation and neurocognitive implications. *Brain Behav Immun* 45:1–12. <https://doi.org/10.1016/j.bbi.2014.10.008>
- Hurst RS, Hajós M, Raggenbass M et al (2005) A novel positive allosteric modulator of the alpha7 neuronal nicotinic acetylcholine receptor: in vitro and in vivo characterization. *J Neurosci* 25:4396–4405. <https://doi.org/10.1523/JNEUROSCI.5269-04.2005>
- INSIGHT START Study Group, Lundgren JD, Babiker AG et al (2015) Initiation of antiretroviral therapy in early asymptomatic HIV infection. *N Engl J Med* 373:795–807. <https://doi.org/10.1056/NEJMoal506816>
- Johnston GR, Webster NR (2009) Cytokines and the immunomodulatory function of the vagus nerve. *Br J Anaesth* 102:453–462. <https://doi.org/10.1093/bja/aep037>
- Joint United Nations Programme on HIV/AIDS (UNAIDS) (2013) Global report: UNAIDS report on global AIDS epidemic 2013
- Kaiser S, Wonnacott S (2000) alpha-bungarotoxin-sensitive nicotinic receptors indirectly modulate [^3H]dopamine release in rat striatal slices via glutamate release. *Mol Pharmacol* 58:312–318
- Kalashnyk OM, Gergalova GL, Komisarenko SV, Skok MV (2012) Intracellular localization of nicotinic acetylcholine receptors in human cell lines. *Life Sci* 91:1033–1037. <https://doi.org/10.1016/j.lfs.2012.02.005>
- Kanmogne GD, Schall K, Leibhart J et al (2007) HIV-1 gp120 compromises blood–brain barrier integrity and enhance monocyte

- migration across blood–brain barrier: implication for viral neuropathogenesis. *J Cereb Blood Flow Metab* 27:123–134. <https://doi.org/10.1038/sj.jcbfm.9600330>
- Kaul M, Garden GA, Lipton SA (2001) Pathways to neuronal injury and apoptosis in HIV-associated dementia. *Nature* 410:988–994. <https://doi.org/10.1038/35073667>
- Kawashima K, Fujii T, Moriwaki Y, Misawa H (2012) Critical roles of acetylcholine and the muscarinic and nicotinic acetylcholine receptors in the regulation of immune function. *Life Sci* 91:1027–1032. <https://doi.org/10.1016/j.lfs.2012.05.006>
- Kawashima K, Fujii T, Moriwaki Y et al (2015) Non-neuronal cholinergic system in regulation of immune function with a focus on $\alpha 7$ nAChRs. *Int Immunopharmacol* 29:127–134. <https://doi.org/10.1016/j.intimp.2015.04.015>
- Koutsilieri E, Czub S, Scheller C et al (2000) Brain choline acetyltransferase reduction in SIV infection. An index of early dementia? *Neuroreport* 11:2391–2393
- Koval LM, Yu Lykhmus O, Omelchenko DM et al (2009) The role of alpha7 nicotinic acetylcholine receptors in B lymphocyte activation. *Ukr Biokhimicheskii Zhurnal* 81:5–11
- Lindgren S, Stewenius J, Sjölund K et al (1993) Autonomic vagal nerve dysfunction in patients with ulcerative colitis. *Scand J Gastroenterol* 28:638–642
- Lindl KA, Marks DR, Kolson DL, Jordan-Sciutto KL (2010) HIV-associated neurocognitive disorder: pathogenesis and therapeutic opportunities. *J Neuroimmune Pharmacol* 5:294–309. <https://doi.org/10.1007/s11481-010-9205-z>
- Liu QH, Williams DA, McManus C et al (2000) HIV-1 gp120 and chemokines activate ion channels in primary macrophages through CCR5 and CXCR4 stimulation. *Proc Natl Acad Sci USA* 97:4832–4837. <https://doi.org/10.1073/pnas.090521697>
- Liu H, Liu J, Liang S, Xiong H (2013) Plasma gelsolin protects HIV-1 gp120-induced neuronal injury via voltage-gated K^+ channel Kv2.1. *Mol Cell Neurosci* 57:73–82
- Liu L, Yu J, Li L et al (2017) Alpha7 nicotinic acetylcholine receptor is required for amyloid pathology in brain endothelial cells induced by Glycoprotein 120, methamphetamine and nicotine. *Sci Rep* 7:40467. <https://doi.org/10.1038/srep40467>
- Loram LC, Harrison JA, Chao L et al (2010) Intrathecal injection of an alpha seven nicotinic acetylcholine receptor agonist attenuates gp120-induced mechanical allodynia and spinal pro-inflammatory cytokine profiles in rats. *Brain Behav Immun* 24:959–967. <https://doi.org/10.1016/j.bbi.2010.03.008>
- Mansvelder HD, Mertz M, Role LW (2009) Nicotinic modulation of synaptic transmission and plasticity in cortico-limbic circuits. *Semin Cell Dev Biol* 20:432–440. <https://doi.org/10.1016/j.semdb.2009.01.007>
- Martin JH (2003) *Neuroanatomy: text and atlas*, 3rd edn. McGraw-Hill Companies, New York
- Maschke M, Kastrup O, Esser S et al (2000) Incidence and prevalence of neurological disorders associated with HIV since the introduction of highly active antiretroviral therapy (HAART). *J Neurol Neurosurg Psychiatry* 69:376–380
- McGuire JL, Barrett JS, Vezina HE et al (2014) Adjuvant therapies for HIV-associated neurocognitive disorders. *Ann Clin Transl Neurol* 1:938–952. <https://doi.org/10.1002/acn3.131>
- Meneses G, Bautista M, Florentino A et al (2016) Electric stimulation of the vagus nerve reduced mouse neuroinflammation induced by lipopolysaccharide. *J Inflamm* 13:33. <https://doi.org/10.1186/s12950-016-0140-5>
- Mihalak KB, Carroll FI, Luetje CW (2006) Varenicline is a partial agonist at alpha4beta2 and a full agonist at alpha7 neuronal nicotinic receptors. *Mol Pharmacol* 70:801–805. <https://doi.org/10.1124/mol.106.025130>
- Nizri E, Irony-Tur-Sinai M, Lory O et al (2009) Activation of the cholinergic anti-inflammatory system by nicotine attenuates neuroinflammation via suppression of Th1 and Th17 responses. *J Immunol Baltim Md* 190:6681–6688. <https://doi.org/10.4049/jimmunol.0902212>
- Norman GJ, Morris JS, Karelina K et al (2011) Cardiopulmonary arrest and resuscitation disrupts cholinergic anti-inflammatory processes: a role for cholinergic $\alpha 7$ nicotinic receptors. *J Neurosci* 31:3446–3452. <https://doi.org/10.1523/JNEUROSCI.4558-10.2011>
- Ochoa ELM, Lasalde-Dominicci J (2007) Cognitive deficits in schizophrenia: focus on neuronal nicotinic acetylcholine receptors and smoking. *Cell Mol Neurobiol* 27:609–639. <https://doi.org/10.1007/s10571-007-9149-x>
- Park-Wyllie LY, Antoniou T (2003) Concurrent use of bupropion with CYP2B6 inhibitors, nelfinavir, ritonavir and efavirenz: a case series. *AIDS Lond Engl* 17:638–640. <https://doi.org/10.1097/01.aids.0000050846.71999.ff>
- Paterson D, Nordberg A (2000) Neuronal nicotinic receptors in the human brain. *Prog Neurobiol* 61:75–111
- Pedrol-Clotet E, Deig-Comerma E, Ribell-Bachs M et al (2006) Bupropion use for smoking cessation in HIV-infected patients receiving antiretroviral therapy. *Enfermedades Infecc Microbiol Clínica* 24:509–511
- Pohanka M (2012) Alpha7 nicotinic acetylcholine receptor is a target in pharmacology and toxicology. *Int J Mol Sci* 13:2219–2238. <https://doi.org/10.3390/ijms13022219>
- Price RW, Perry SW (1994) *HIV, AIDS, and the brain*. Raven Press Ltd, New York
- Raber J, Toggas SM, Lee S et al (1996) Central nervous system expression of HIV-1 Gp120 activates the hypothalamic-pituitary-adrenal axis: evidence for involvement of NMDA receptors and nitric oxide synthase. *Virology* 226:362–373. <https://doi.org/10.1006/viro.1996.0664>
- Ramos FM, Delgado-Vélez M, Ortiz ÁL et al (2015) Expression of CHRFAM7A and CHRNA7 in neuronal cells and postmortem brain of HIV-infected patients: considerations for HIV-associated neurocognitive disorder. *J Neurovirol*. <https://doi.org/10.1007/s13365-015-0401-8>
- Rangwala F, Drisdell RC, Rakhilin S et al (1997) Neuronal alpha-bungarotoxin receptors differ structurally from other nicotinic acetylcholine receptors. *J Neurosci* 17:8201–8212
- Rock RB, Gekker G, Aravalli RN et al (2008) Potentiation of HIV-1 expression in microglial cells by nicotine: involvement of transforming growth factor-beta 1. *J Neuroimmune Pharmacol* 3:143–149. <https://doi.org/10.1007/s11481-007-9098-7>
- Sahdeo S, Wallace T, Hirakawa R et al (2014) Characterization of RO5126946, a Novel $\alpha 7$ nicotinic acetylcholine receptor-positive allosteric modulator. *J Pharmacol Exp Ther* 350:455–468. <https://doi.org/10.1124/jpet.113.210963>
- Saylor D, Dickens AM, Sacktor N et al (2016) HIV-associated neurocognitive disorder—pathogenesis and prospects for treatment. *Nat Rev Neurol* 12:234–248. <https://doi.org/10.1038/nrneurol.2016.27>
- Shirley DK, Kaner RJ, Glesby MJ (2013) Effects of smoking on non-AIDS-related morbidity in HIV-infected patients. *Clin Infect Dis* 57:275–282. <https://doi.org/10.1093/cid/cit207>
- Shytle RD, Mori T, Townsend K et al (2004) Cholinergic modulation of microglial activation by alpha 7 nicotinic receptors. *J Neurochem* 89:337–343. <https://doi.org/10.1046/j.1471-4159.2004.02347.x>
- Simioni S, Cavassini M, Annoni J-M et al (2010) Cognitive dysfunction in HIV patients despite long-standing suppression of viremia. *AIDS Lond Engl* 24:1243–1250. <https://doi.org/10.1097/QAD.0b013e3283354a7b>
- Skok MV (2009) Editorial: To channel or not to channel? Functioning of nicotinic acetylcholine receptors in leukocytes. *J Leukoc Biol* 86:1–3. <https://doi.org/10.1189/JLB.0209106>

- Skok M, Grailhe R, Agenes F, Changeux J-P (2006) The role of nicotinic acetylcholine receptors in lymphocyte development. *J Neuroimmunol* 171:86–98. <https://doi.org/10.1016/j.jneuroim.2005.09.011>
- Solomon DA, Sax PE (2015) Current state and limitations of daily oral therapy for treatment. *Curr Opin HIV AIDS* 10:219–225. <https://doi.org/10.1097/COH.000000000000165>
- Spudich S (2013) HIV and neurocognitive dysfunction. *Curr HIV/AIDS Rep* 10:235–243. <https://doi.org/10.1007/s11904-013-0171-y>
- Su X, Lee JW, Matthay ZA et al (2007) Activation of the alpha7 nAChR reduces acid-induced acute lung injury in mice and rats. *Am J Respir Cell Mol Biol* 37:186–192. <https://doi.org/10.1165/rcmb.2006-0240OC>
- Su X, Feng X, Terrando N et al (2012) Dysfunction of inflammation-resolving pathways is associated with exaggerated postoperative cognitive decline in a rat model of the metabolic syndrome. *Mol Med Camb Mass* 18:1481–1490. <https://doi.org/10.2119/molmed.2012.00351>
- Suzuki T, Hide I, Matsubara A et al (2006) Microglial alpha7 nicotinic acetylcholine receptors drive a phospholipase C/IP3 pathway and modulate the cell activation toward a neuroprotective role. *J Neurosci Res* 83:1461–1470. <https://doi.org/10.1002/jnr.20850>
- Swingler S, Mann AM, Zhou J et al (2007) Apoptotic killing of HIV-1-infected macrophages is subverted by the viral envelope glycoprotein. *PLoS Pathog* 3:1281–1290. <https://doi.org/10.1371/journal.ppat.0030134>
- Tavazzi E, Morrison D, Sullivan P et al (2014) Brain inflammation is a common feature of HIV-infected patients without HIV encephalitis or productive brain infection. *Curr HIV Res* 12:97–110
- Tewari M, Monika null, Varghse RK et al (2015) Astrocytes mediate HIV-1 Tat-induced neuronal damage via ligand-gated ion channel P2×7R. *J Neurochem* 132:464–476. <https://doi.org/10.1111/jnc.12953>
- Thomsen MS, Hansen HH, Timmerman DB, Mikkelsen JD (2010) Cognitive improvement by activation of alpha7 nicotinic acetylcholine receptors: from animal models to human pathophysiology. *Curr Pharm Des* 16:323–343
- Tracey KJ (2002) The inflammatory reflex. *Nature* 420:853–859. <https://doi.org/10.1038/nature01321>
- Tracey KJ (2007) Physiology and immunology of the cholinergic anti-inflammatory pathway. *J Clin Invest* 117:289–296. <https://doi.org/10.1172/JCI30555>
- Tracey KJ (2009) Reflex control of immunity. *Nat Rev Immunol* 9:418–428. <https://doi.org/10.1038/nri2566>
- UNAIDS (2017) Fact sheet - Latest statistics on the status of the AIDS epidemic. <https://www.unaids.org/en/resources/fact-sheet>. Accessed 12 July 2018
- Valdés-Ferrer SI, Crispín JC, Belaunzarán PF et al (2009) Acetylcholine-esterase inhibitor pyridostigmine decreases T cell overactivation in patients infected by HIV. *AIDS Res Hum Retrovir* 25:749–755. <https://doi.org/10.1089/aid.2008.0257>
- Valdés-Ferrer SI, Crispín JC, Belaunzarán-Zamudio PF et al (2017) Add-on pyridostigmine enhances CD4⁺ T-cell recovery in HIV-1-infected immunological non-responders: a proof-of-concept study. *Front Immunol* 8:1301. <https://doi.org/10.3389/fimmu.2017.01301>
- Van Westerloo DJ, Giebelen IAJ, Florquin S et al (2005) The cholinergic anti-inflammatory pathway regulates the host response during septic peritonitis. *J Infect Dis* 191:2138–2148. <https://doi.org/10.1086/430323>
- Van Westerloo DJ, Giebelen IA, Florquin S et al (2006) The vagus nerve and nicotinic receptors modulate experimental pancreatitis severity in mice. *Gastroenterology* 130:1822–1830. <https://doi.org/10.1053/j.gastro.2006.02.022>
- Van Maanen MA, Papke RL, Koopman FA et al (2015) Two novel α7 nicotinic acetylcholine receptor ligands: in vitro properties and their efficacy in collagen-induced arthritis in mice. *PLoS ONE* 10:e0116227. <https://doi.org/10.1371/journal.pone.0116227>
- Van der Zanden EP, Hilbers FW, Verseijden C et al (2012) Nicotinic acetylcholine receptor expression and susceptibility to cholinergic immunomodulation in human monocytes of smoking individuals. *Neuroimmunomodulation* 19:255–265. <https://doi.org/10.1159/000335185>
- Vázquez-Palacios G, Bonilla-Jaime H (2004) Nicotine acetylcholine receptors and neuropsychiatric disorders. *Rev Neurol* 39:1146–1160
- Wang H, Yu M, Ochani M et al (2003) Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation. *Nature* 421:384–388. <https://doi.org/10.1038/nature01339>
- Wang D, Zhou R, Yao Y (2009) Role of cholinergic anti-inflammatory pathway in regulating host response and its interventional strategy for inflammatory diseases. *Chin J Traumatol* 12:355–364
- Wang D, Zhou R, Yao Y et al (2010) Stimulation of α7 nicotinic acetylcholine receptor by nicotine increases suppressive capacity of naturally occurring CD4⁺ CD25⁺ regulatory T cells in mice in vitro. *J Pharmacol Exp Ther* 335:553–561. <https://doi.org/10.1124/jpet.110.169961>
- Woods SP, Moore DJ, Weber E, Grant I (2009) Cognitive neuropsychology of HIV-associated neurocognitive disorders. *Neuropsychol Rev* 19:152–168. <https://doi.org/10.1007/s11065-009-9102-5>
- Wu J, George AA, Schroeder KM et al (2004) Electrophysiological, pharmacological, and molecular evidence for alpha7-nicotinic acetylcholine receptors in rat midbrain dopamine neurons. *J Pharmacol Exp Ther* 311:80–91. <https://doi.org/10.1124/jpet.104.070417>
- Yoshikawa H, Kurokawa M, Ozaki N et al (2006) Nicotine inhibits the production of proinflammatory mediators in human monocytes by suppression of I-kappaB phosphorylation and nuclear factor-kappaB transcriptional activity through nicotinic acetylcholine receptor alpha7. *Clin Exp Immunol* 146:116–123. <https://doi.org/10.1111/j.1365-2249.2006.03169.x>
- Zhang B, Yu J-Y, Liu L-Q et al (2015) Alpha7 nicotinic acetylcholine receptor is required for blood-brain barrier injury-related CNS disorders caused by *Cryptococcus neoformans* and HIV-1 associated comorbidity factors. *BMC Infect Dis* 15:352. <https://doi.org/10.1186/s12879-015-1075-9>