

# Ethanol Effect on BK Channels is Modulated by Magnesium

Héctor G. Marrero, Steven N. Treistman, and José R. Lemos

**Background:** Alcoholics have been reported to have reduced levels of magnesium in both their extracellular and intracellular compartments. Calcium-dependent potassium channels (BK) are known to be one of ethanol (EtOH)'s better known molecular targets.

**Methods:** Using outside-out patches from hippocampal neuronal cultures, we examined the consequences of altered intracellular  $Mg^{2+}$  on the effects that EtOH has on BK channels.

**Results:** We find that the effect of EtOH is bimodally influenced by the  $Mg^{2+}$  concentration on the cytoplasmic side. More specifically, when internal  $Mg^{2+}$  concentrations are  $\leq 200 \mu M$ , EtOH decreases BK activity, whereas it increases activity when  $Mg^{2+}$  is at 1 mM. Similar results are obtained when using patches from HEK cells expressing only the  $\alpha$ -subunit of BK. When patches are made with the actin destabilizer cytochalasin D present on the cytoplasmic side, the potentiation caused by EtOH becomes independent of the  $Mg^{2+}$  concentration. Furthermore, in the presence of the actin stabilizer phalloidin, EtOH causes inhibition even at  $Mg^{2+}$  concentrations of 1 mM.

**Conclusions:** Internal  $Mg^{2+}$  can modulate the EtOH effects on BK channels only when there is an intact, internal actin interaction with the channel, as is found at synapses. We propose that the EtOH-induced decrease in cytoplasmic  $Mg^{2+}$  observed in frequent/chronic drinkers would decrease EtOH's actions on synaptic (e.g., actin-bound) BK channels, producing a form of molecular tolerance.

**Key Words:** Ethanol, Magnesium, BK Channel.

**I**N THIS STUDY, we examined the consequences of altered  $Mg^{2+}$  concentrations on one of ethanol (EtOH)'s better known molecular targets, the calcium-dependent potassium channel (BK). BK activity is potentiated by EtOH, and this modulation is dependent on the intracellular  $Ca^{2+}$  concentration (Liu et al., 2008, 2013; Yuan et al., 2011). More specifically, EtOH increases the activity of the channel when internal  $Ca^{2+}$  concentrations (1 to 30  $\mu M$ ) are at physiological levels (Feinberg-Zadek and Treistman, 2007; Feinberg-Zadek et al., 2008; Liu et al., 2008). EtOH's potentiation of BK channel activity would lead to a stronger action potential after hyperpolarization, thus reducing firing frequency. The final effect would be a less responsive neuronal system (Gruss et al., 2001).

The  $Ca^{2+}$  dependence of EtOH's effects on BK channels has been well documented (see Mulholland et al., 2009). Cytoplasmic  $Ca^{2+}$  is thought to be the only cation necessary for the EtOH potentiation of these channels (Liu et al., 2008). This is concluded despite the fact that  $Mg^{2+}$  affects BK activity (Cui et al., 2009; Horrigan and Ma, 2008; Hu et al., 2003, 2007; Yang et al., 2008; Zhang et al., 2001) in a similar manner as does  $Ca^{2+}$ .  $Mg^{2+}$ , at physiological levels,

increases BK channel open probability and is not only competitive with  $Ca^{2+}$  at some binding sites but also has independent  $Mg^{2+}$ -specific binding sites at the c-terminus of BK (Chen et al., 2011; Cui, 2010; Cui et al., 2009; Latorre and Brauchi, 2006; Yang et al., 2008; see Fig. 6). Thus, it would be reasonable to expect that  $Mg^{2+}$  could also modulate EtOH's effects on BK channels.

The cytoskeleton influences the activity of BK channels. It is well known that the presence of polymerized actin (as is found at synapses; Frotscher et al., 2014; Gordon-Weeks and Fournier, 2014; Loebrich, 2014; Mori et al., 2014) causes a decrease in the activity of some types of BK channels (Brainard et al., 2005; O'Malley et al., 2005) perhaps through interaction with sites at the c-terminus (Tian et al., 2006). Moreover, actin polymerization is dependent on the  $Mg^{2+}$  concentration (Galińska-Rakoczy et al., 2009; Hild et al., 2010). Interestingly, EtOH affects cytoskeletal actin filament integrity (Allansson et al., 2001; Loureiro et al., 2011; Offenhäuser et al., 2006; Popp and Dertien, 2008; Romero et al., 2010; Szabo et al., 2007). Importantly, alcohol consumption and/or exposure results in the depletion of  $Mg^{2+}$  levels in serum (Altura and Altura, 1999; Brown et al., 2002; Poikolainen and Alho, 2008; Romani, 2008) as well as cellular internal  $Mg^{2+}$  (Babu et al., 1999; Li et al., 2001; Romani, 2008). These are compelling reasons to explore whether cytoplasmic  $Mg^{2+}$  and actin can modulate the effects of EtOH on ionic channels.

We report here that EtOH's effects on BK channel activity are influenced by internal  $Mg^{2+}$  and that these effects are dependent on the integrity of the internal actin structure. Furthermore, this influence is observed at  $Ca^{2+}$  concentrations reported to be high enough (with respect to magnesium

From the Institute of Neurobiology UPR-MSU (HGM, SNT), San Juan, Puerto Rico; and Department of Microbiology and Physiological Systems (JRL), University of Massachusetts Medical School, Worcester, Massachusetts.

Received for publication May 5, 2015; accepted June 17, 2015.

Reprint requests: Héctor G. Marrero, PhD, Institute of Neurobiology UPR-MSU, 201 Blvd. del Valle, San Juan, PR 00901; Tel.: +1-787-721-4149; Fax: +1-787-725-3804, E-mail: hector.marrero3@upr.edu

Copyright © 2015 by the Research Society on Alcoholism.

DOI: 10.1111/acer.12821



concentrations) to avoid  $\text{Ca}^{2+}$ -binding site competition (Shi and Cui, 2001; see Zhang et al., 2001). Thus, the influence of  $\text{Mg}^{2+}$  should be considered to be independent of other modulators (e.g.,  $\text{Ca}^{2+}$ ) of EtOH's effects on BK channels. We also discuss the possible regulation of BK channels by the well-known " $\text{Mg}^{2+}$  deficit" syndrome resulting from chronic EtOH consumption, where the decrease in potentiation of channel activity by acute EtOH could be a form of tolerance resulting from these changes in internal  $\text{Mg}^{2+}$  levels.

## MATERIALS AND METHODS

### Preparations

Hippocampal CA1 neurons from embryonic rats were cultured for 7 to 14 days before use. Neurons from these cultures are known to only express  $\alpha$  and  $\alpha$ - $\beta 4$  subunits of BK channels (Misonou et al., 2006; Piwonska et al., 2008). The culture media were then slowly changed to external solutions, usually requiring 20 to 30 volume equivalents. The procedure would last about 5 minutes and did not show any visible sign of damage or trauma to the cells, which could be patched within approximately 4 hours after the solution exchange.

### Solutions

In all cases, the external (bath) solution used was (in mM): 130.5 K-gluconate, 14.5 KCl, 10 HEPES, 3 NaCl, 2 HEGTA, pH 7.35 (using KOH) with  $\text{MgCl}_2$  and  $\text{CaCl}_2$  added to make free concentrations of 1 mM and 30  $\mu\text{M}$ , respectively. The amounts used for the free ion concentrations were calculated using the program WEB-MAXC STANDARD (online program, 2009 version, Stanford University), with ionic strength at 0.16, and at 20°C. Internal (pipette) solutions were similar to external, except the amounts of  $\text{MgCl}_2$  and  $\text{CaCl}_2$  were prepared to achieve different free ion concentrations, as indicated. All tests for voltage shifts ( $V_{1/2}$ , see below) were made with internal 30  $\mu\text{M}$  free  $\text{Ca}^{2+}$ , to minimize possible competition with free  $\text{Mg}^{2+}$  on the high-affinity  $\text{Ca}^{2+}$  sites (Chen et al., 2011; Cui, 2010; Lee and Cui, 2010) of BK channels. One millimolar free external  $\text{Mg}^{2+}$  was used because, first, the approach of studying internal  $\text{Mg}^{2+}$  effects would be simplified by changing only its internal concentration. Second, it was thought that one important question to answer would be what would happen if conditions mimic the internal  $\text{Mg}^{2+}$  depletion caused by EtOH (see Discussion). Finally, it is much easier to obtain stable outside-out configurations when patches are made with high external/bath  $\text{Mg}^{2+}$ .

The agents used were dimethyl sulfoxide (DMSO), cytochalasin D (both from Sigma-Aldrich, St. Louis, MO), and phalloidin (Invitrogen, Eugene, OR). All were diluted to their final concentrations in the internal solutions. Dilutions of cytochalasin D and phalloidin would yield a final DMSO concentration of 0.1% (v/v).

The "moderate" EtOH concentration range of 20 to 25 mM is the lower-limit of "intoxicating" in terms of human and animal consumption (see Eckardt et al., 1998; Harris and Mihic, 2004) and has been previously used for studies in EtOH effects on decreasing internal cellular  $\text{Mg}^{2+}$  (see, e.g., Babu et al., 1999; Li et al., 2001). Thus, in all cases here, EtOH was applied directly to the bath to make a final concentration of 20 to 25 mM.

### Patching Procedure

Borosilicate glass pipettes were made with a resistance of 7 to 10 Mega-Ohms (M $\Omega$ ). Pipettes were first tip-filled with the particular internal solution to be tested without any agents, and then back-filled with the same solutions plus agents, whenever such were tested. High-resistance on-cell patches were created (>2 Giga-Ohms:

G $\Omega$ ) followed by the whole-cell, and finally outside-out configurations. To improve the stability of the seals, in most cases, the patches were left unperturbed for at least 5 minutes after obtaining the outside-out configuration.

### Data Gathering

Pulse protocols and data recordings were made using a HEKA EPC10 amplifier controlled with PATCHMASTER program (all by HEKA Elektronik, Dr. Schulze GmbH, Lambrecht/Pfalz, Germany). A series of criteria were followed to include recordings in our analysis. First, in accordance with what is known from hippocampal BK channels (Gong et al., 2001), patches with channel openings with conductances below 220 pS were not considered. Second, voltage-dependent activity was monitored through the use of voltage ramps, from which a range of voltage-dependent activity was determined (minimum to maximum activity), and an initial visual estimate of the voltage for NPo = 0.5 could be made. One of 2 protocols was followed once the patch was determined as stable.

*Tests Using Changes in Specific Value of NPo.* Patches were held at a holding potential where NPo was estimated to be about 0.5 (from the voltage ramps). A series of square pulses were then given, in increments of 1 mV and of 5 seconds duration, from -5 mV to +5 mV from that holding potential, until a recording yielded a control NPo = 0.5 to 0.6.

*NPo as a Function of Voltage.* Recordings were made using square pulses, each of 3 to 5 seconds duration, in increments of 5 to 10 mV from the lowest to slightly above the highest used in the voltage ramps (see Data Gathering). The higher limit was obtained by setting the voltage to where no closed states could be distinguished. The criteria for the duration of each test took into account that a minimum of 3 consecutive stable sets should be taken just prior to EtOH application and that the total time lapsed for all 3 should not be more than 2 to 5 minutes. This is of particular importance when considering that acute EtOH effects on the BK channel are usually reported to occur within the first 2 to 5 minutes after drug application (e.g., see Martin et al., 2008). Thus, recordings in the presence of EtOH would be limited to the next 2 to 5 minutes after EtOH application.

*Analysis.* All-points amplitude histograms were obtained from recordings where discrete channel openings, as well as closed states, could be distinguished. From there, the NPo was calculated using

$$\text{NPo} = \frac{\sum_i iA_i}{\sum_i A_i},$$

where  $N$  is the number of open levels,  $i$  the level number ( $i = 0$  is the closed state), and  $A_i$  is the area of a Gaussian fit to the  $i$ th level in the all-points histogram. The NPo for each voltage, from at least 3 consecutive sets (see above), was averaged and either included in a set of  $n$ 's (0.5 to 0.6 NPo bar graphs) or plotted versus applied voltage.

*$V_{1/2}$  Measurements.* When  $V_{1/2}$  was measured, the criterion for using each plot was to have at least 2 points in the voltage ranges where saturation of NPo was observed. Fits of the NPo-versus-voltage plots were made using

$$N_m \text{PO} = \frac{N_m}{1 + e^{k(V_{1/2} - V)}}$$

(IgorPro, v. 6.1; WaveMetrics, Inc., Portland, OR), where  $N_m$  is the maximum number of levels larger than zero. The fits corresponds to a Boltzmann function multiplied by  $N_m$ , with  $k$  having the usual value  $zF/KT$ . From this, the EtOH-induced shifts in  $V_{1/2}$  (from controls) were determined.

**Statistics.** In order to avoid tolerance to EtOH, only 1 cell per dish was used. Thus,  $n$  represents the number of patches = number of cells = number of culture dishes used. For the  $V_{1/2}$ -versus-voltage tests, a minimum of  $n = 5$  per case was used to comply with the basic binomial distribution requirements (Shapiro–Wilk test). Two-way analyses of variance (ANOVAs) were performed for the tests with specific values of NPo (i.e., = 0.5 to 0.6, fixed variables were  $X_1$ : with/without EtOH and  $X_2$ :  $[Mg^{2+}]$ ) as well as for the results from NPo versus voltage (fixed variables were  $X_1$ :  $[Mg^{2+}] = 200 \mu M/[Mg^{2+}] = 1 \text{ mM}$  and  $X_2$ : treatments). Tukey tests were made post hoc the ANOVA tests, to determine specific pair differences. Significant differences in all tests were considered when  $p < 0.05$  and are reported following the indication of the type of test.

## RESULTS

### *EtOH-Induced Changes in BK Activity are Regulated by Cytoplasmic Magnesium*

It is well known that EtOH changes the open probability (NPo) of BK channels. We worked at NPo values near 0.5 to 0.6 as controls (see Methods) to detect either positive or negative EtOH-induced changes. Measurements were conducted with the cytoplasmic side of the patch exposed to buffered free  $Ca^{2+}$  at  $30 \mu M$  but different concentrations of buffered free  $Mg^{2+}$  (Fig. 1). Surprisingly, significant EtOH-induced decreases in NPo were observed with cytoplasmic-side  $Mg^{2+}$  concentrations  $\leq 200 \mu M$  (Fig. 1). In contrast, significant EtOH-induced increases were observed in the presence of  $1 \text{ mM } Mg^{2+}$ . There was no change in NPo at  $400 \mu M$ . Similar results (data not shown) are observed with  $5 \mu M$  cytoplasmic  $Ca^{2+}$ . These results demonstrate that (i) there is a  $Mg^{2+}$ -dependent EtOH effect on the activity of BK channels in outside-out patches and (ii) the effect is not affected by  $Ca^{2+}$  at the physiological concentrations used here for both ions (Shi and Cui, 2001).


### *Magnesium-Dependent EtOH Effects versus Voltage*

BK channels are regulated by both  $Ca^{2+}$  and voltage. Therefore, BK channel NPo was recorded as a function of voltages that included observed minima and maxima of NPo. Unless associated with a change in conductance or channel density, changes in specific values of NPo (at particular voltages) are the result of shifts in the  $V_{1/2}$  of such NPo-versus-voltage curves. Thus, to obtain a more clear and complete description of the NPo-versus-voltage characteristics, the  $V_{1/2}$  shift was used to determine EtOH-induced potentiation (negative shift) or depression (positive shift) of BK activity. A total of  $200 \mu M Mg^{2+}$  was used as a comparison with  $1 \text{ mM } Mg^{2+}$ , both at  $30 \mu M$  internal  $Ca^{2+}$ .


Figure 2 shows an example of EtOH-induced increases in BK activity when internal  $Mg^{2+}$  is at  $1 \text{ mM}$ . The EtOH-induced increase in channel activity is an effect that is within the voltage ranges that include the observed minima and maxima of NPo (Fig. 2D). In contrast, EtOH induces a decrease in BK activity when internal  $Mg^{2+}$  is at  $200 \mu M$  (Fig. 3D). A similar behavior was observed when patches from HEK cells, expressing only the BK- $\alpha$  subunit, were used, indicating that this phenomenon could be a property of  $\alpha$ -BK channels without other subunits (Fig. 4). Thus, the EtOH-induced effects on the activity of the BK channels can be observed throughout the physiological voltage range.

### *The Magnesium-Dependent EtOH Effects on BK Channels are Dependent on Associated Actin Filaments*


In contrast to our results, potentiation by EtOH was previously reported to be independent of internal  $Mg^{2+}$  (e.g., see Liu et al., 2008). As actin filaments interact with BK channels (Tian et al., 2006), an explanation for this difference could be the preservation of actin filaments in the outside-out patch configuration (Ruknudin et al., 1991), as opposed to other configurations (where internal components are expected to be scarce or absent), such as inside-out (e.g., Liu et al., 2008, 2013; Yuan et al., 2011) or artificial bilayers




**Fig. 1.** Magnesium-dependent ethanol (EtOH) influence on calcium-dependent potassium channels (BK) from hippocampal neurons. Shown are EtOH-induced changes from initial control NPo values near 0.5 to 0.6. Significant EtOH-induced NPo reduction is observed at  $\leq 200 \mu M$  free  $Mg^{2+}$  suggesting that the transition from potentiation to depression might be mostly dependent on the  $Mg^{2+}$  concentration. These results show that the  $200 \mu M$  magnesium cases are a reasonable and consistent choice (see Methods) for comparison with the  $1 \text{ mM } Mg^{2+}$  cases in the more rigorous tests that followed. Data shown are for internal  $30 \mu M$  free  $Ca^{2+}$ . Similar effects seen using  $5 \mu M$  free  $Ca^{2+}$  (data not shown). White-filled bars are control and gray-filled are +EtOH. Error bars are SE. The brackets under the horizontal axis indicate significant difference ( $t$ -test's  $p < 0.016$ ) between a paired-case with the same free  $Mg^{2+}$  (bottom row), with  $n$  indicated below each control/+EtOH pair (row above magnesium concentrations). The control data were targeted to NPo values between 0.5 and 0.6. As voltages for controls were different throughout the experiments, the +EtOH NPo is reported as that obtained at the same voltage as that of the corresponding control. Following ANOVA of all data ( $\pm$  EtOH vs.  $[Mg^{2+}]$ ,  $p < 0.0001$ ), Tukey tests indicated a significant difference limit at  $200 \mu M Mg^{2+}$  when +EtOH is compared to its control (at specific  $[Mg^{2+}]$  for Control vs. +EtOH;  $100 \mu M$ :  $p < 0.0145$ ,  $200 \mu M$ :  $p < 0.0485$ ,  $400 \mu M$ :  $p = 1$ ,  $600 \mu M$ :  $p > 0.9997$ ,  $1 \text{ mM}$ :  $p < 0.003$ ).



**Fig. 2.** Example of ethanol (EtOH)-induced potentiation of calcium-dependent potassium channels (BK) obtained from hippocampal neurons. Internal solution contained  $30 \mu\text{M}$  free  $\text{Ca}^{2+}$  and  $1 \text{ mM}$  free  $\text{Mg}^{2+}$ . (A) Addition of 20 to 25 mM EtOH causes an increase in activity. All-points histograms (sample of **B** = control and **C** = +EtOH) were used to obtain plots of NPo versus applied voltage (**D**). In **D**, fits made of these plots (dashed black = control, dashed gray = +EtOH) indicate that EtOH causes a leftward voltage shift. The examples given in **A** (and for **B** and **C**) were from records taken at a voltage of  $-5 \text{ mV}$ . In **B** and **C**, raw histograms are shown as black traces, Gaussian fits as gray traces, and composite of Gaussian fits as dashed gray traces. Similar effects seen using  $5 \mu\text{M}$  free  $\text{Ca}^{2+}$  (data not shown).



**Fig. 3.** Example of ethanol (EtOH)-induced depression of calcium-dependent potassium channels (BK) obtained from hippocampal neurons. Internal solution contained  $30 \mu\text{M}$  free  $\text{Ca}^{2+}$  and  $200 \mu\text{M}$  free  $\text{Mg}^{2+}$ . (A) Addition of EtOH causes a decrease in activity. All-points histograms (sample of **B** = control and **C** = +EtOH) were used to obtain plots of NPo versus applied voltage (**D**). In **D**, fits made for these plots (dashed black = control, dashed gray = +EtOH) indicate that EtOH causes a rightward voltage shift. The examples given in **A** (and for **B** and **C**) were from records taken at a voltage of  $-30 \text{ mV}$ . All other plot details are as explained in Fig. 2.



**Fig. 4.** Examples of ethanol (EtOH)-induced changes on calcium-dependent potassium channels (BK) activity from HEK cells expressing only the BK- $\alpha$  subunit. With 1 mM internal (cytoplasmic) free  $Mg^{2+}$ , addition of EtOH causes an increase in activity (**A**). This activity increase is seen in the plots of NPo-versus-applied voltage (**B**) as a leftward shift in  $V_{1/2}$  (dashed black = control, dashed gray = +EtOH). There were 4 trials ( $n = 4$ ) made for this case, giving an average  $V_{1/2}$  shift of ( $mV \pm SE$ )  $-12.14 \pm 2.37$ . The example given in **A** was from records taken at  $-20$  mV (contained in **B**). With  $200 \mu M$  internal free  $Mg^{2+}$ , addition of EtOH causes a decrease in activity (**C**), also seen in the plots of NPo-versus-applied voltage (**D**) as a rightward shift in  $V_{1/2}$  (dashed black = control, dashed gray = +EtOH). There were 4 trials ( $n = 4$ ) made for this case, giving an average  $V_{1/2}$  shifts of ( $mV \pm SE$ )  $11.27 \pm 0.77$ , significantly different from the  $V_{1/2}$  shift obtained with 1 mM free  $Mg^{2+}$  ( $t$ -test  $p < 0.0002$ ). The example given in **C** was from records taken at  $-60$  mV (contained in **D**). Internal solution for all tests had  $30 \mu M$  free  $Ca^{2+}$ .

(e.g., Crowley et al., 2003; Pau et al., 2011; Yuan et al., 2011).


To test this, agents that would either disrupt ( $10 \mu M$  cytochalasin D) or stabilize ( $10 \mu M$  phalloidin) actin were used when testing for EtOH effects with both high and low internal  $Mg^{2+}$ . The possible effect of DMSO was also considered, as it is the vehicle solvent for phalloidin and cytochalasin, but it yielded no significant effects on the EtOH-induced  $V_{1/2}$  shifts in NPo. The average control  $V_{1/2}$  values with  $200 \mu M$  free  $Mg^{2+}$  ( $=mV \pm SE$ ): normal =  $-7.12 \pm 4.22$  ( $n = 15$ ), +DMSO =  $-26.98 \pm 5.55$  ( $n = 10$ ), +Phalloidin =  $-14.30 \pm 5.67$  ( $n = 11$ ), and +Cytochalasin D =  $-7.98 \pm 3.40$  ( $n = 15$ ); for 1 mM free  $Mg^{2+}$ : normal =  $1.02 \pm 3.90$  ( $n = 8$ ), +DMSO =  $-17.98 \pm 6.68$  ( $n = 7$ ), +Phalloidin =  $-8.28 \pm 7.44$  ( $n = 9$ ), and +Cytochalasin D =  $10.18 \pm 7.19$  ( $n = 7$ ). The effects found for EtOH on  $V_{1/2}$  values did not depend on their initial (or control)  $V_{1/2}$  values. A summary of the EtOH-induced effect  $V_{1/2}$  shifts in NPo is given in Fig. 5.

Interestingly, when  $Mg^{2+}$  is  $200 \mu M$ , then EtOH decreases BK activity unless actin is disrupted by cytochalasin D. In contrast, EtOH potentiates the activity with 1 mM  $Mg^{2+}$  unless actin is stabilized by phalloidin.

Thus, when actin filaments are destabilized, then EtOH induces a potentiation of BK channel activity, with either high or low internal  $Mg^{2+}$ . In contrast, when actin filaments are stabilized by phalloidin, then EtOH depresses BK channel activity, regardless of the  $Mg^{2+}$  concentration. It is clear that the EtOH-induced changes of BK channel activity are determined by internal  $Mg^{2+}$ , but that this reliance is contingent on the integrity of the actin filaments associated with the channel.

## DISCUSSION

Here, we report for the first time that, in outside-out patches,  $Mg^{2+}$  alters, in a concentration-dependent manner, the effects of EtOH on BK channel activity. Previously, experimental conditions used to test for EtOH effects on BK channels involved the use of cell-attached techniques, inside-out patches, or reconstituted channels in artificial bilayers. In these, internal  $Mg^{2+}$  is not controlled (e.g., cell attached), or the participation of internal components has been minimized (e.g., inside-out) or eliminated (e.g., artificial bilayers). In our case, internal cellular components should be retained with the outside-out patch configuration (Ruknudin et al., 1991). Therefore, the conditions obtained for single channel recordings better approximate those at the surface of intact cells. Furthermore, we used  $[Mg^{2+}]/[Ca^{2+}]$  ratios that are lower than that reported for  $Mg^{2+}$ - $Ca^{2+}$  competition for the high-affinity  $Ca^{2+}$  binding site (see Shi and Cui, 2001). Thus, it appears that any influence of actin on the high-affinity binding site is minimal, and therefore, we only considered influences on low-affinity binding sites for  $Mg^{2+}$  (see Fig. 6).



**Fig. 5.** Summary of ethanol (EtOH)-induced  $V_{1/2}$  shifts from hippocampal neurons. Potentiation corresponds to negative  $V_{1/2}$  shifts (bars below horizontal axis), while depression corresponds to positive  $V_{1/2}$  shifts (bars above horizontal axis). Dimethyl sulfoxide (DMSO) was at 0.1%, also present at this concentration when phalloidin (**Phal**; 10  $\mu$ M) or cytochalasin D (**CD**; 10  $\mu$ M) were used. The *ns* are given in the lowest row. EtOH-induced potentiation is present with 1 mM free  $Mg^{2+}$ , while depression is induced with 200  $\mu$ M free  $Mg^{2+}$ . The presence of CD causes a persistent EtOH-induced potentiation, while Phal causes a persistent depression. Following ANOVA of all data (treatment versus  $[Mg^{2+}]$ ,  $p < 0.0001$ ), Tukey tests indicated that within the 200  $\mu$ M  $Mg^{2+}$ , the results with CD were significantly different from the other 3 (largest  $p < 0.0110$ ), whereas at 1 mM  $Mg^{2+}$ , it was Phal which was significantly different from the other 3 (largest  $p < 0.030$ ). The “\*” indicates the case which was significantly different from the others with the same  $Mg^{2+}$  concentration. Black capped lines are standard errors (SE).

### The EtOH-Induced Shift in $V_{1/2}$ and Cytoskeletal Components


EtOH-induced activity changes are contingent on the cytoplasmic  $Mg^{2+}$  concentrations. The  $Mg^{2+}$ -dependent EtOH effect on BK channels is only evident if they are still associated with actin filaments. Furthermore, these effects demonstrate, for the first time, that  $Mg^{2+}$  is a possible modulator of EtOH's effects. However, there is not yet enough information to determine whether the effects are a direct influence on the BK channel's  $Mg^{2+}$  binding sites, or on the ion's effect on actin polymerization, or some mixture of both (see Fig. 6). In accordance with this model, actin filaments are known to be stabilized by increasing  $Mg^{2+}$  (Galińska-Rakoczy et al., 2009; Hild et al., 2010). Thus, EtOH-induced depression would have been expected for high  $Mg^{2+}$  concentrations. That this is not the case indicates that the causes for the  $Mg^{2+}$ -dependent EtOH effect cannot be explained by the direct effect of  $Mg^{2+}$  on the actin filaments. Still, the  $Mg^{2+}$ -EtOH interaction appears to be a result of the presence of actin filaments or actin components, such as found at synapses (Frotscher et al., 2014; Gordon-Weeks and Fournier, 2014; Loebrich, 2014; Mori et al., 2014).

### Hypothesis on the Dependency of $Mg^{2+}$ Action on Actin

Figure 6 is a model that includes the participation of the c-terminus of the BK channel in the EtOH- $Mg^{2+}$ -actin effect. In this model, actin modulates the voltage-dependent openings of the BK channel, such modulation being in turn dependent on the  $Mg^{2+}$  concentration. The modulation of open probability is simplified as directly associated to move-

ment of the BK c-terminus (Fig. 6). The model considers only binding sites for  $Mg^{2+}$  (which are at the c-terminus), and actin, as well as possible EtOH interaction sites at the channel protein and bilayer sides.

The main premise of the model is that the channel's activity would depend on the stabilization of the c-terminus with respect to the membrane, with the highest open probability occurring when the terminus is closest to the main transmembrane segments of the  $\alpha$  subunit. There is already precedence for this feature of the model, as others have suggested that a similar re-conformation of the c-terminus is associated with the open state of K-channels (Cui et al., 2009; Jiang et al., 2002). Thus, any condition that increases the *possibility* of having the c-terminus closer to the membrane would also increase the channel's open probability. The model includes certain assumptions that are focused on effects stabilizing the position of the c-terminus: (i)  $Mg^{2+}$  binding to its site in the channel would favor stabilization of the c-terminus near the membrane, (ii) actin filament binding would hinder movement of the c-terminus, and (iii) EtOH binding would favor c-terminus movement toward or away from the membrane, depending on the c-terminus position previous to EtOH application. Taken together, the presence of bound, polymerized actin would stabilize the conditions present previous to EtOH application. Notice that although the model does not include the effect of  $Mg^{2+}$  on the actin filaments' stability, it is still consistent with what is known about  $Mg^{2+}$ -dependent actin polymerization. Namely, under low  $Mg^{2+}$ , it is expected that the effect of “less polymerized” actin would result in a more negative control's  $V_{1/2}$ , but having an opposite effect with higher  $Mg^{2+}$ .



**Fig. 6.** Model of the mechanism of action of ethanol (EtOH)-induced changes in calcium-dependent potassium channels (BK) activity. An inherent mobility of the c-terminus (C-T) part of the channel is assumed and associated with BK activity. The basis for modulation of activity would be the C-T's interaction with the  $\alpha$  trans membrane sections (and bilayer). Thus, one main feature of the model is the proposition that anything that alters the probability of the C-T being closer to the bilayer would also increase the probability of channel openings (depicted by thickness of the white arrows inside the trans membrane section " $\alpha$ ", at top). The high/low  $Mg^{2+}$  (left/right of vertical dashed line) would correspondingly increase/decrease the open probability. Actin filaments would stabilize either state, thus determining the modulation of the system by EtOH. Thus, if the C-T was stabilized close to the bilayer (by actin, bottom drawing at left of vertical dashed line), addition of EtOH will further stabilize (or bring closer) the C-T. On the other hand, if the C-T were actin-stabilized away from the bilayer, addition of EtOH would further stabilize it away from the bilayer (bottom drawing at right of vertical dashed line). See Discussion for more details.

### Another View of EtOH Tolerance

One major consequence of alcohol consumption and/or exposure is the depletion (down to 200 to 400  $\mu$ M from 1 mM) of cellular internal  $Mg^{2+}$  (see Babu et al., 1999; Li et al., 2001; Romani, 2008). This depletion can occur even with small amounts of EtOH and as early as the first few minutes after ingestion or exposure (Babu et al., 1999). The long-term consequences are usually highly symptomatic, and therapeutic strategies for  $Mg^{2+}$  replenishment have been used as remedies for alcoholism (Poikolainen and Alho, 2008). We now hypothesize that the BK channel is affected by the  $Mg^{2+}$  depletion caused by EtOH. Namely, that the EtOH-induced reduction of intracellular  $Mg^{2+}$  would differentially alter BK channel activity in cellular areas where BK channels are highly associated with actin filaments. More explicitly, synapses,

where there is high actin polymerization, would show a decrease in BK activity when internal  $Mg^{2+}$  is reduced by EtOH. Although at the cellular level, this would have various consequences; at the systemic level, the overall effect can be interpreted as another mechanism for tolerance in habitual (or chronic) EtOH drinkers. Given that while in occasional drinkers, the  $Mg^{2+}$  levels would recover shortly after cessation of EtOH consumption, in habitual drinkers, the  $Mg^{2+}$  levels would remain low for extended periods (Eckardt et al., 1998; Poikolainen and Alho, 2008; Torres et al., 2009).

Thus, we hypothesize that in occasional drinkers, EtOH induces potentiation of BK activity, but that there would be no potentiation (i.e., tolerance) at synaptic (i.e., actin-bound) BK channels, because of the decrease in cytoplasmic  $Mg^{2+}$  observed in frequent/chronic drinkers.

In summary, EtOH effects on BK channel are modulated by the concentration of internal (cytoplasmic)  $Mg^{2+}$ . This effect is dependent upon the integrity of actin elements associated with BK channels such as those at synaptic membranes. Thus, for most alcoholics, the low levels of internal  $Mg^{2+}$  would contribute to tolerance of the EtOH-induced potentiation of BK channel activity.

## ACKNOWLEDGMENTS

This work was supported by grants from NIH NS29470 to JRL, NIH R01AA018038-06 and R01A8380110, both to SNT, and funds supporting HGM provided by NIH 5P20GM103642 (Centers of Biomedical Research Excellence). We would like to give special thanks to Dr. Cristina M. Velazquez for helpful discussions and in the editing of the manuscript. No conflict of interest is associated with the authors and the contents of this manuscript, and thus none to declare.

## REFERENCES

- Allansson L, Khatibi S, Olsson T, Hansson E (2001) Acute ethanol exposure induces  $[Ca^{2+}]_i$  transients, cell swelling and transformation of actin cytoskeleton in astroglial primary cultures. *J Neurochem* 76:472–479.
- Altura BM, Altura BT (1999) Association of alcohol in brain injury, headaches and stroke with brain-tissue and serum levels of ionized magnesium: a review of recent-findings and mechanisms of action. *Alcohol* 19:111–130.
- Babu AN, Cheng TP, Zhang A, Altura BT, Altura BM (1999) Low concentrations of ethanol deplete type-2 astrocytes of intracellular free magnesium. *Brain Res Bull* 50:59–62.
- Brainard AM, Miller AJ, Martens JR, England SK (2005) Maxi-K channels localize to caveolae in human myometrium: a role for an actin-channel-caveolin complex in the regulation of myometrial smooth muscle  $K^+$  current. *Am J Physiol Cell Physiol* 289:C49–C57.
- Brown RA, Ilg KJ, Chen AF, Ren J (2002) Dietary  $Mg(2+)$  supplementation restores impaired vasoactive responses in isolated rat aorta induced by chronic ethanol consumption. *Eur J Pharmacol* 442:241–250.
- Chen R-S, Geng Y, Magleby KL (2011)  $Mg^{2+}$  binding to open and closed states can activate BK channels provided that the voltage sensors are elevated. *J Gen Physiol* 138:593–607.
- Crowley JJ, Treistman SN, Dopico AM (2003) Cholesterol antagonizes ethanol potentiation of human brain BKCa channels reconstituted into phospholipid bilayers. *Mol Pharmacol* 64:365–372.
- Cui J (2010) BK-type calcium-activated potassium channels: coupling of metal ions and voltage sensing. *J Physiol* 588(Pt 23):4651–4658.
- Cui J, Yang H, Lee US (2009) Molecular mechanisms of BK channel activation. *Cell Mol Life Sci* 66:852–875.
- Eckardt MJ, File SE, Gessa GL, Grant KA, Guerri C, Hoffman PL, Kalant H, Koob GF, Li TK, Tabakoff B (1998) Effects of moderate alcohol consumption on the central nervous system. *Alcohol Clin Exp Res* 22:998–1040.
- Feinberg-Zadek PL, Martin G, Treistman SN (2008) BK channel subunit composition modulates molecular tolerance to ethanol. *Alcohol Clin Exp Res* 32:1207–1216.
- Feinberg-Zadek PL, Treistman SN (2007) Beta-subunits are important modulators of the acute response to alcohol in human BK channels. *Alcohol Clin Exp Res* 31:737–744.
- Frotscher M, Studer D, Graber W, Chai X, Nestel S, Zhao S (2014) Fine structure of synapses on dendritic spines. *Front Neuroanat* 8:94.
- Galińska-Rakoczy A, Wawro B, Strzelecka-Gołaszewska H (2009) New aspects of the spontaneous polymerization of actin in the presence of salts. *J Mol Biol* 387:869–882.
- Gong LW, Gao TM, Huang H, Tong Z (2001) Properties of large conductance calcium-activated potassium channels in pyramidal neurons from the hippocampal CA1 region of adult rats. *Jpn J Physiol* 51:725–731.
- Gordon-Weeks PR, Fournier AE (2014) Neuronal cytoskeleton in synaptic plasticity and regeneration. *J Neurochem* 129:206–212.
- Gruss M, Henrich M, König P, Hempelmann G, Vogel W, Scholz A (2001) Ethanol reduces excitability in a subgroup of primary sensory neurons by activation of BK(Ca) channels. *Eur J Neurosci* 14:1246–1256.
- Harris RA, Mihic SJ (2004) Alcohol and inhibitory receptors: unexpected specificity from a nonspecific drug. *PNAS* 101:2–3.
- Hild G, Bugyi B, Nyitrai M (2010) Conformational dynamics of actin: effectors and implications for biological function. *Cytoskeleton* 67:609–629.
- Horrigan FT, Ma Z (2008)  $Mg^{2+}$  enhances voltage sensor/gate coupling in BK channels. *J Gen Physiol* 131:13–32.
- Hu L, Shi J, Ma Z, Krishnamoorthy G, Sieling F, Zhang G, Horrigan FT, Cui J (2003) Participation of the S4 voltage sensor in the  $Mg^{2+}$ -dependent activation of large conductance (BK)  $K^+$  channels. *Proc Natl Acad Sci USA* 100:10488–10493.
- Hu L, Yang H, Shi J, Cui J (2007) Effects of multiple metal binding sites on calcium and magnesium-dependent activation of BK channels. *J Gen Physiol* 127:35–49.
- Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R (2002) Crystal structure and mechanism of a calcium-gated potassium channel. *Nature* 417:515–522.
- Latorre R, Brauchi S (2006) Large conductance  $Ca^{2+}$ -activated  $K^+$  (BK) channel: activation by  $Ca^{2+}$  and voltage. *Biol Res* 39:385–401.
- Lee US, Cui J (2010) BK channel activation: structural and functional insights. *Trends Neurosci* 33:415–423.
- Li W, Zheng T, Babu AN, Altura BT, Gupta RK, Altura BM (2001) Importance of magnesium ions in development of tolerance to ethanol: studies on cultured cerebral vascular smooth muscle cells, type-2 astrocytes and intact rat brain. *Brain Res Bull* 56:153–158.
- Liu J, Bukiya AN, Kuntamallappanavar G, Singh AK, Dopico AM (2013) Distinct sensitivity of slo1 channel proteins to ethanol. *Mol Pharmacol* 83:235–244.
- Liu J, Vaithianathan T, Manivannan K, Parrill A, Dopico AM (2008) Ethanol modulates BKCa channels by acting as an adjuvant of calcium. *Mol Pharmacol* 74:628–640.
- Loeblich S (2014) The role of F-actin in modulating Clathrin-mediated endocytosis: lessons from neurons in health and neuropsychiatric disorder. *Commun Integr Biol* 7:e28740.
- Loureiro SO, Heimfarth L, Reis K, Wild L, Andrade C, Guma FT, Gonçalves CA, Pessoa-Pureur R (2011) Acute ethanol exposure disrupts actin cytoskeleton and generates reactive oxygen species in c6 cells. *Toxicol In Vitro* 25:28–36.
- Martin GE, Hendrickson LM, Penta KL, Friesen RM, Pietrzykowski AZ, Tapper AR, Treistman SN (2008) Identification of a BK channel auxiliary protein controlling molecular and behavioral tolerance to alcohol. *Proc Natl Acad Sci USA* 105:17543–17548.
- Misonou H, Menegola M, Buchwalder L, Park EW, Meredith A, Rhodes KJ, Aldrich RW, Trimmer JS (2006) Immunolocalization of the  $Ca^{2+}$ -activated  $K^+$  channel Slo1 in axons and nerve terminals of mammalian brain and cultured neurons. *J Comp Neurol* 496:289–302.
- Mori M, Rikitake Y, Mandai K, Takai Y (2014) Roles of nectins and nectin-like molecules in the nervous system. *Adv Neurobiol* 8:91–116.
- Mulholland PJ, Hopf FW, Bukiya AN, Martin GE, Liu J, Dopico AM, Bonci A, Treistman SN, Chandler LJ (2009) Sizing up ethanol-induced plasticity: the role of small and large conductance calcium-activated potassium channels. *Alcohol Clin Exp Res* 33:1125–1135.
- Offenhäuser N, Castelletti D, Mapelli L, Soppo BE, Regondi MC, Rossi P, D'Angelo E, Frassoni C, Amadeo A, Tocchetti A, Pozzi B, Disanza A, Guarneri D, Betsholtz C, Scita G, Heberlein U, Di Fiore PP (2006) In-



- creased ethanol resistance and consumption in Eps8 knockout mice correlates with altered actin dynamics. *Cell* 127:213–226.
- O'Malley D, Irving AJ, Harvey J (2005) Leptin-induced dynamic changes in the actin cytoskeleton mediate the activation and synaptic clustering of BK channels. *FASEB J* 19:1917–1919. Available at: <http://www.fasebj.org/cgi/doi/10.1096/fj.05-4166fje>. Accessed August 12, 2014.
- Pau VPT, Smith FJ, Taylor AB, Parfenova LV, Samakai E, Callaghan MM, Abarca-Heidemann K, Hart PJ, Rothberg BS (2011) Structure and function of multiple Ca<sup>2+</sup>-binding sites in a K<sup>+</sup> channel regulator of K<sup>+</sup> conductance (RCK) domain. *Proc Natl Acad Sci USA* 108:17684–17689.
- Piwonska M, Wilczek E, Szewczyk A, Wilczynski GM (2008) Differential distribution of Ca<sup>2+</sup>-activated potassium channel  $\beta$ 4 subunit in rat brain: immunolocalization in neuronal mitochondria. *Neuroscience* 153:446–460.
- Poikolainen K, Alho H (2008) Magnesium treatment in alcoholics: a randomized clinical trial. *Subst Abuse Treat Prev Policy* 3:1.
- Popp RL, Dertien JS (2008) Actin depolymerization contributes to ethanol inhibition of NMDA receptors in primary cultured cerebellar granule cells. *Alcohol* 42:525–539.
- Romani AM (2008) Magnesium homeostasis and alcohol consumption. *Magnes Res* 21:197–204.
- Romero AM, Esteban-Pretel G, Marín MP, Ponsoda X, Ballestín R, Canales JJ, Renau-Piqueras J (2010) Chronic ethanol exposure alters the levels, assembly, and cellular organization of the actin cytoskeleton and microtubules in hippocampal neurons in primary culture. *Toxicol Sci* 118:602–612.
- Ruknudin A, Song MJ, Sachs F (1991) The ultrastructure of patch-clamped membranes: a study using high voltage electron microscopy. *J Cell Biol* 112:125–134.
- Shi J, Cui J (2001) Intracellular Mg<sup>2+</sup> enhances the function of BK-type Ca<sup>2+</sup>-activated K(+) channels. *J Gen Physiol* 118:589–606.
- Szabo G, Dolganiuc A, Dai Q, Pruett SB (2007) TLR4, ethanol, and lipid rafts: a new mechanism of ethanol action with implications for other receptor-mediated effects. *J Immunol* 178:1243–1249.
- Tian L, Chen L, McClafferty H, Sailer CA, Ruth P, Knaus H-G, Shipston MJ (2006) A noncanonical SH3 domain binding motif links BK channels to the actin cytoskeleton via the SH3 adapter cortactin. *FASEB J* 20:E2046–E2056.
- Torres LM, Cefaratti C, Berti-Mattera L, Romani A (2009) Delayed restoration of Mg<sup>2+</sup> content and transport in liver cells following ethanol withdrawal. *Am J Physiol Gastrointest Liver Physiol* 297:G621–G631.
- Yang H, Shi J, Zhang G, Yang J, Delaloye K, Cui J (2008) Activation of Slo1 BK channels by Mg<sup>2+</sup> coordinated between the voltage sensor and the RCK1 domains. *Nat Struct Mol Biol* 15:1152–1159.
- Yuan C, Chen M, Covey DF, Johnston LJ, Treistman SN (2011) Cholesterol tuning of BK ethanol response is enantioselective, and is a function of accompanying lipids. *PLoS ONE* 6:e27572.
- Zhang X, Solaro CR, Lingle CJ (2001) Allosteric regulation of BK channel gating by Ca<sup>2+</sup> and Mg<sup>2+</sup> through a nonselective, low affinity divalent cation site. *J Gen Physiol* 118:607–635.